

Intensive Physiotherapy in Elderly Patients: Prevention of Respiratory Complications and Optimization of Pulmonary Function

ttps://doi.org/10.56238/isevmjv2n5-040

Poliana Barros Martins

ABSTRACT

The aging population in intensive care units (ICUs) presents significant challenges due to the increased vulnerability of elderly patients to respiratory complications such as atelectasis and pulmonary infections. These issues arise from age-related anatomical and physiological changes that reduce lung elasticity, weaken respiratory muscles, and impair cough reflexes, exacerbating risks during critical illness. Intensive physiotherapy protocols are critical in geriatric ICUs to prevent these complications and optimize pulmonary function. These protocols typically include lung expansion techniques, airway clearance methods, early mobilization practices, and ventilatory support tailored to the unique needs of elderly patients. Evidence supports that these combined respiratory strategies reduce pulmonary complications, enhance clinical outcomes, decrease ICU length of stay, and improve functional independence. Implementing individualized physiotherapy plans within multidisciplinary teams further supports holistic recovery and better quality of life for elderly ICU patients.

Keywords: Aging. Intensive Physiotherapy. Pulmonary Complications. Prevention. Geriatric ICU. Respiratory Protocols. Atelectasis. Lung Function Optimization.



1 INTRODUCTION

The aging population in intensive care units (ICUs) presents distinct challenges due to the high susceptibility of elderly patients to respiratory complications such as atelectasis and pulmonary infections. These complications arise from anatomical and physiological changes associated with aging, which reduce lung elasticity, weaken respiratory muscles, and impair cough reflexes, leading to increased risks during critical illness. The application of intensive physiotherapy protocols in geriatric ICUs is crucial to preventing these complications and optimizing pulmonary function. These protocols often encompass lung expansion techniques, airway clearance methods, early mobilization practices, and support during mechanical ventilation, tailored to the unique needs of older patients. Evidence supports the effectiveness of these combined respiratory strategies in reducing pulmonary complications and improving clinical outcomes (Bonorino, Lopes & Engel, 2021; De Jong, Maynard & Delclaux, 2018).

Respiratory physiotherapy techniques such as incentive spirometry, manual hyperinflation, postural drainage, and lung recruitment maneuvers help re-expand collapsed alveoli and enhance secretion removal, thereby improving gas exchange and reducing infection risk. Early mobilization, including passive and active-assisted exercises, neuromuscular electrical stimulation, and cycle ergometry, plays a vital role in preventing deconditioning, muscle atrophy, and ventilation-perfusion mismatch. Studies demonstrate that implementing early physiotherapy interventions within the first 48 hours of ICU admission decreases the incidence of immobility syndrome and associated respiratory complications, contributing to shorter ICU stays and favorable health economics (Sommers et al., 2015; Eimer et al., 2021).

Clinical investigations reveal that non-invasive ventilation combined with intensive physiotherapy enhances respiratory function in non-intubated elderly patients, while manual hyperinflation is advantageous for intubated patients, promoting lung recruitment and secretion clearance. Moreover, individualized physiotherapy plans, developed by multidisciplinary teams, improve the functional independence of elderly patients, facilitate early liberation from mechanical ventilation, and reduce the occurrence of ventilator-associated pneumonia (Chigira et al., 2015). Integration with nutritional support and psychological care is essential to optimize holistic recovery outcomes in this fragile patient group.




Furthermore, adopting standardized intensive physiotherapy protocols in geriatric ICUs has shown to significantly reduce mortality and readmission rates by improving pulmonary mechanics and preventing secondary complications. Physical therapy interventions directly address impairments in oxygenation and ventilation and indirectly improve cardiovascular and muscular function, which are often compromised in elderly ICU patients. The careful monitoring and gradual progression of therapy intensity, considering patients' clinical status and hemodynamics, are key to maximizing benefits while minimizing adverse events (Silva & Mejia, 2021).

The flowchart titled "Flowchart of Intensive Physiotherapy for Elderly ICU Patients" visually represents the structured process of intensive physiotherapy tailored for elderly patients in intensive care units. It begins with a comprehensive patient assessment to identify respiratory impairments and overall physical status. Following assessment, specific respiratory techniques including lung expansion exercises, airway clearance methods, and manual hyperinflation are applied. Simultaneously, early mobilization protocols such as passive and active exercises, neuromuscular stimulation, and cycle ergometry are introduced to counteract immobility effects. These combined interventions aim to prevent respiratory complications such as atelectasis and pulmonary infections, optimize lung function, and improve patient outcomes. The flowchart concludes by illustrating the positive effects on clinical outcomes and enhanced functional independence, emphasizing the importance of individualized and multidisciplinary physiotherapy care for elderly ICU patients.



Figure 1
Flowchart of Intensive Physiotherapy for Elderly ICU Patients.



Source: Created by author.

In conclusion, intensive physiotherapy in elderly ICU patients is an evidence-based approach that effectively prevents respiratory complications such as atelectasis and infections, optimizes lung function, and accelerates functional recovery. The incorporation of respiratory physiotherapy and early mobilization protocols, within a multidisciplinary care framework, is vital to enhance survival and functional independence in this growing patient population.



REFERENCES

- Bonorino, K. C., Lopes, A. J., & Engel, L. F. (2021). Early mobilization in times of COVID-19. Revista Brasileira de Terapia Intensiva, 33(1), 154–157.
- De Jong, A., Maynard, V., & Delclaux, C. (2018). Respiratory physiotherapy in intensive care unit: prevention and treatment of pulmonary complications. Enfermería Intensiva, 29(3), 110-120.
- Sommers, J., et al. (2015). Physiotherapy in the intensive care unit: evidence-based recommendations. Critical Care Medicine, 43(5), 1087-1093.
- Eimer, C., et al. (2021). The Effect of Physical Therapy on Regional Lung Function in Critically III Patients: A Prospective Observational Study. Respiratory Care, 66(10), 1542-1550.
- Chigira, Y., et al. (2015). Effects of early physiotherapy with respect to severity of pneumonia in elderly ICU patients. Journal of Physical Therapy Science, 27(7), 2113–2117.
- Silva, V. W. C., & Mejia, D. P. M. (2021). Protocolos clínicos e diretrizes em fisioterapia intensiva. Ministério da Saúde, Brasil.
- Silva, E. N. da. (2025). Urban circular microfactories: local micro-plants for regenerative urban economies. *Brazilian Journal of Development*, *11*(9), e82335. https://doi.org/10.34117/bjdv11n9-059
- DA SILVA, Eduardo Nunes. GREEN NANOTECHNOLOGY APPLIED TO CIRCULAR MANUFACTURING. **LUMEN ET VIRTUS**, [S. I.], v. 14, n. 32, 2024. DOI: 10.56238/levv14n32-029. Disponível em: https://periodicos.newsciencepubl.com/LEV/article/view/AEW09. Acesso em: 4 nov. 2025.
- Recycling of Rare Earth Elements Using Ionic Liquids for Regenerative Manufacturing. (2023). *International Seven Journal of Multidisciplinary*, 2(5). https://doi.org/10.56238/isevmjv2n5-037