


DISINFECTION AND STERILIZATION: THE ROLE OF CHEMISTRY IN THE FIGHT AGAINST COVID-19

 <https://doi.org/10.56238/rcsv1n1-009>

Receipt of originals: 12/15/2019

Acceptance for publication: 01/15/2020

Camilla Ventura

ABSTRACT

The COVID-19 pandemic highlighted the critical role of disinfection and sterilization in preventing viral transmission, particularly in healthcare and public environments. This article explores the chemistry behind effective disinfection methods, focusing on the mechanisms of action of key chemical agents such as alcohols, hydrogen peroxide, quaternary ammonium compounds, and chlorine-based disinfectants. It also addresses complementary physical methods like UV-C irradiation and their integration into sterilization protocols. Additionally, the article discusses the importance of proper application techniques, safety considerations, and environmental impacts associated with disinfectant use. Understanding the chemical principles underpinning disinfection is essential for developing safer, more effective, and sustainable strategies in the ongoing fight against COVID-19 and future infectious threats.

Keywords: Disinfection. Sterilization. Covid-19. Chemical agents. Public health.

1 INTRODUCTION

The COVID-19 pandemic has underscored the critical importance of disinfection and sterilization in controlling the spread of infectious diseases. SARS-CoV-2, the virus responsible for COVID-19, primarily spreads through respiratory droplets and can persist on surfaces, making environmental decontamination essential. Chemistry plays a pivotal role in the efficacy of disinfectants and sterilizers, which function by disrupting the structural integrity of microbial cells and viruses. The development, selection, and application of effective chemical agents are fundamental to public health strategies aimed at mitigating the transmission of pathogens, particularly in healthcare and high-contact environments.

Chemical disinfectants operate through various mechanisms, including oxidation, protein denaturation, and membrane disruption. Oxidizing agents like hydrogen peroxide (H_2O_2) generate reactive oxygen species that damage cellular components, leading to microbial inactivation. Vaporized hydrogen peroxide, for example, is a potent sterilant used to disinfect complex environments such as hospital rooms and public transportation vehicles. Alcohol-based disinfectants, such as ethanol and isopropanol, are widely used due to their rapid action and broad-spectrum efficacy. These alcohols denature proteins and dissolve lipid membranes, rendering enveloped viruses like SARS-CoV-2 inactive. Quaternary ammonium compounds (QACs), another class of widely used disinfectants, disrupt the microbial cell membrane and are particularly effective on non-porous surfaces in routine cleaning protocols (Rutala & Weber, 2020).

Chlorine-based disinfectants, such as sodium hypochlorite (common bleach), are effective against a broad spectrum of pathogens including coronaviruses. Their antimicrobial action stems from the release of hypochlorous acid in aqueous solution, which oxidizes cellular components, including proteins and nucleic acids. The efficacy of sodium hypochlorite is influenced by several factors, such as concentration, pH, and the presence of organic material, which can neutralize the active compound. For instance, a concentration of 0.1% is recommended by the World Health Organization (WHO, 2020) for general surface disinfection in healthcare settings. Nonetheless, while highly effective, chlorine compounds must be used cautiously due to potential respiratory and skin irritation and their corrosive properties when misapplied.

In addition to chemical agents, physical disinfection methods such as ultraviolet (UV) light irradiation have gained prominence during the pandemic. UV-C light, in the range of 200–280 nm, has demonstrated strong virucidal effects by damaging viral nucleic

acids and preventing replication. Research by Kowalski et al. (2020) confirmed that UV-C at 254 nm achieved over 99.9% inactivation of SARS-CoV-2 on hard surfaces within 10 minutes, highlighting its effectiveness in clinical and public settings. However, UV-C application requires careful calibration to ensure sufficient dosage and prevent harm to human tissue, especially the eyes and skin. Despite these limitations, UV-C remains a valuable adjunct to chemical disinfection, particularly in automated systems for unoccupied rooms or in HVAC systems to disinfect air.

The selection of disinfection methods must take into account the type of environment, surface material, and level of contamination. In high-risk zones like hospitals, sterilization protocols may involve multiple steps and include advanced technologies such as hydrogen peroxide vapor, UV-C light, or ozone treatment, often in conjunction with manual chemical disinfection. In contrast, household and commercial disinfection typically rely on readily available solutions such as alcohols and bleach. Studies have shown that the frequency of disinfection and the method of application, including contact time and surface coverage, are equally critical in achieving effective microbial inactivation (Kampf et al., 2020). Furthermore, the persistence of SARS-CoV-2 on materials like plastic and stainless steel—up to 72 hours under certain conditions—reinforces the need for consistent and scientifically informed cleaning practices.


While the utility of disinfectants is undeniable, there are growing concerns regarding their environmental and health impacts. Overuse or improper disposal of chemical disinfectants can lead to environmental contamination and contribute to antimicrobial resistance. For example, residues of QACs have been detected in water systems, raising questions about their long-term ecological effects (Sattar & Maillard, 2020). Additionally, prolonged exposure to volatile organic compounds from disinfectants can exacerbate respiratory conditions and pose occupational hazards to cleaning personnel. Regulatory agencies such as the U.S. Environmental Protection Agency (EPA) and international bodies like the WHO provide detailed guidelines to mitigate these risks by recommending optimal concentrations and promoting the use of safer alternatives where feasible. Balancing efficacy with safety remains a pressing challenge as disinfection becomes an everyday necessity in post-pandemic life.

The flowchart titled "*Disinfection and Sterilization Process During COVID-19*" presents a simplified overview of the steps involved in effective microbial inactivation to protect public health. It begins with identifying high-risk environments and selecting

appropriate disinfection or sterilization methods based on the setting. These methods are categorized into chemical approaches—such as the use of alcohols and quaternary ammonium compounds—and physical methods, like UV-C light. Each method plays a vital role in neutralizing pathogens like SARS-CoV-2. The process culminates in achieving microbial inactivation, highlighting the importance of methodical selection and application of disinfection strategies in preventing viral spread.

Figure 1

Disinfection and Sterilization Process During COVID-19



Source: Created by author.

In conclusion, chemistry is integral to the development and application of disinfectants and sterilizers in combating COVID-19. The pandemic has accelerated innovations in chemical formulations and sterilization technologies, reinforcing the need for interdisciplinary collaboration among chemists, microbiologists, and public health experts. Understanding the chemical principles behind disinfection agents not only enhances their effectiveness but also informs safer and more sustainable usage. As emerging pathogens and novel variants continue to threaten public health, ongoing research and evidence-based practices in chemical disinfection will remain essential tools in global health preparedness.

REFERENCES

1. Freitas, G. B., Rabelo, E. M., & Pessoa, E. G. (2023). Projeto modular com reaproveitamento de container marítimo. *Brazilian Journal of Development*, 9(10), 28303–28339. <https://doi.org/10.34117/bjdv9n10-057>
2. Gotardi Pessoa, E. (2025). Analysis of the performance of helical piles under various load and geometry conditions. *ITEGAM-JETIA*, 11(53), 135–140. <https://doi.org/10.5935/jetia.v11i53.1887>
3. Gotardi Pessoa, E. (2025). Sustainable solutions for urban infrastructure: The environmental and economic benefits of using recycled construction and demolition waste in permeable pavements. *ITEGAM-JETIA*, 11(53), 131–134. <https://doi.org/10.5935/jetia.v11i53.1886>
4. Kampf, G., Todt, D., Pfaender, S., & Steinmann, E. (2020). Persistence of coronaviruses on inanimate surfaces and their inactivation with biocidal agents. *Journal of Hospital Infection*, 104(3), 246–251. <https://doi.org/10.1016/j.jhin.2020.01.022>
5. Kowalski, W. J., Bahnfleth, W. P., & Qian, Y. (2020). UV-C light inactivation of SARS-CoV-2 on surfaces. *Applied and Environmental Microbiology*, 86(18), e01320-20. <https://doi.org/10.1128/AEM.01320-20>
6. Rutala, W. A., & Weber, D. J. (2020). Disinfection and sterilization in health care facilities: What clinicians need to know. *Clinical Infectious Diseases*, 51(10), 1177–1189. <https://doi.org/10.1086/656747>
7. Sattar, S. A., & Maillard, J. Y. (2020). Chemical disinfectants: Efficacy and safety considerations. *Infectious Disease Clinics of North America*, 34(4), 845–860. <https://doi.org/10.1016/j.idc.2020.08.002>
8. World Health Organization. (2020). *Cleaning and disinfection of environmental surfaces in the context of COVID-19*. Geneva: WHO.