


NFκB1 gene polymorphism increases the risk for occurrence of fatigue and sleep disorder in patients with sporadic PD

O polimorfismo do gene NFκB1 aumenta o risco de ocorrência de fadiga e distúrbio do sono em pacientes com DP esporádica

 <https://doi.org/10.56238/cienciasaudeestuepesv1-086>

Elaine Bandeira Cavalcanti Duarte

Department of Biology, Federal Rural University of Pernambuco (UFRPE), Recife, PE, Brazil

Erinaldo Ubirajara Damasceno dos Santos

Department of Biology, Federal Rural University of Pernambuco (UFRPE), Recife, PE, Brazil

Samantha Amorim Cândido

Department of Biology, Federal Rural University of Pernambuco (UFRPE), Recife, PE, Brazil

Géssica Dayane Cordeiro de Lima

Department of Biology, Federal Rural University of Pernambuco (UFRPE), Recife, PE, Brazil

Amdore Guescel C. Asano

Department of Clinical Medicine, Faculty of Medicine, Federal University of Pernambuco (UFPE), Recife, PE, Brazil

Pro-Parkinson Program of the Clinical Hospital of the Federal University of Pernambuco Recife (HC/UFPE), Recife, PE, Brazil

Nadja Maria Jorge Asano

Department of Clinical Medicine, Faculty of Medicine, Federal University of Pernambuco (UFPE), Recife, PE, Brazil

Pro-Parkinson Program of the Clinical Hospital of the Federal University of Pernambuco Recife (HC/UFPE), Recife, PE, Brazil

Maria de Mascena Diniz Maia

Department of Biology, Federal Rural University of Pernambuco (UFRPE), Recife, PE, Brazil

Paulo Roberto Eleutério de Souza

Department of Biology, Federal Rural University of Pernambuco (UFRPE), Recife, PE, Brazil

ABSTRACT

Objectives The objective of this study was to investigate a possible relationship between genetic variants in the *TNFA* and *NFKB1* genes with the occurrence of fatigue and sleep disorder in patients with Parkinson's disease (PD).

Methods A total of 180 Brazilian patients from the Pro-Parkinson service at the Clinical Hospital of the Federal University of Pernambuco, diagnosed with sporadic PD were enrolled. The sequence variants for *TNFA* and *NFKB1* were determined through the PCR-RFLP technique. Logistic regression was performed to assess the association of polymorphisms with the occurrence of fatigue and sleep disorder in PD patients.

Key findings Multivariate analysis revealed increased risk for the occurrence of fatigue in individuals carrying *NFKB* D/W genotype (Prevalence Ratio: PR=2.091; 95%CI, 1.155-3.784; p=0.015) and D/D genotype (PR= 2.237; 95%CI, 1.078-4.643; p=0.031). In addition, it was observed an increased risk to the occurrence of fatigue in PD patients with sleep disorder (PR= 5.226; 95%CI, 2.934-9.307; p= 0.000). Moreover, the therapy with pramipexole was significantly associated with the development of sleep disorder (p = 0.020).

Conclusion *NFKB1* variants and use of the pramipexole may have a significant influence on increased risk for the occurrence of fatigue and sleep disorder, respectively in a Brazilian PD population.

Keywords: Parkinson's disease, Inflammation, Fatigue, Sleep disorder, Polymorphism.

RESUMO

Objetivos O objetivo deste estudo foi investigar uma possível relação entre as variantes genéticas dos genes *TNFα* e *NFKB1* com a ocorrência de fadiga e distúrbio do sono em pacientes com doença de Parkinson (DP).

Métodos Foram inscritos 180 pacientes brasileiros do serviço Pró-Parkinson do Hospital Clínico da Universidade Federal de Pernambuco, diagnosticados com DP esporádica. As variantes da seqüência para TNF α e NFKB1 foram determinadas através da técnica PCR-RFLP. A regressão logística foi realizada para avaliar a associação de polimorfismos com a ocorrência de fadiga e distúrbio do sono em pacientes com DP.

Principais conclusões A análise multivariada revelou um risco maior para a ocorrência de fadiga em indivíduos portadores do genótipo NFKB D/W (Razão de prevalência: PR=2,091; 95%CI, 1,155-3,784; p=0,015) e genótipo D/D (PR= 2,237; 95%CI, 1,078-4,643; p=0,031). Além disso, foi observado um risco

maior de ocorrência de fadiga em pacientes com distúrbio do sono (PR= 5.226; IC95%, 2.934-9.307; p= 0.000). Além disso, a terapia com pramipexole foi significativamente associada ao desenvolvimento do distúrbio do sono (p = 0,020).

Conclusão As variantes do NFKB1 e o uso do pramipexole podem ter uma influência significativa no aumento do risco de ocorrência de fadiga e distúrbio do sono, respectivamente em uma população de DP brasileira.

Palavras-chave: Doença de Parkinson, Inflamação, Fadiga, Transtorno do sono, Polimorfismo.

1 INTRODUCTION

Parkinson's disease (PD) is considered the second neurodegenerative disease most common in the elderly population over 60 years old [1]. It is characterized by the loss progressive of dopaminergic neurons in the substantia nigra region of the midbrain [2]. Approximately 1% of the worldwide population and 3.3% of the Brazilian population is affected by this disorder [1]. The etiology of PD is still complex, involving immunologic, genetic and environmental factors [3]. The inflammation process may be considered a hallmark of PD since that were reported levels increased of inflammatory molecules in the brain and cerebrospinal fluid of PD patients [4]. Furthermore, Ghosh et al. showed that NF-kB1, activation a transcription regulator of the κ immunoglobulin light chain (IgK), is induced in the CNS of patients with PD[5].

The proinflammatory cytokines released by activated microglia, including tumor necrosis factor α (TNF- α), interleukins IL-6 and IL-1 β , are over-regulated in the brain and cerebrospinal fluid of patients with PD. TNF- α , in particular, can activate neuroinflammation and the molecular mechanisms that lead to neurotoxicity and neuronal death both in vitro and in animal models of PD [6]. In animal models, TNF- α and other proinflammatory cytokines can induce the loss of dopaminergic neurons [7]. Post-mortem studies in patients with PD identified immunoreactive glial cells to TNF- α in the substantia nigra, high levels of TNF- α in nigrostriatal dopaminergic regions and cerebrospinal fluid, and still indicated release of TNF- α from activated glial cells in the substantia nigra [8].

Fatigue and sleep disorder are non-motor symptoms of PD and may cause a great impact on the quality of life of these patients [9]. Some studies on neural-immunological signaling have shown that pro-inflammatory molecules can signal the Central Nervous System (CNS), triggering fatigue and other behavioral changes. These findings raise the hypothesis that fatigue and sleep disorder would be related to the inflammatory process [10,11].

In recent years, genetic mechanisms such as changes in DNA base sequences have begun to be revealed as potential factors in the pathogenesis of PD [12]. Such alterations, known as genetic polymorphisms, play an important role in the regulation of gene expression and in the silencing of repetitive elements of the genome. Thus, genetic variations such as the occurrence of polymorphisms in the NF- κ B1 and TNF- α genes may be able to silence, increase or reduce gene expression, contributing to the pathogenesis and progression of neurodegenerative diseases [13,14].

This way, the purpose of this study was to verify a possible association between variants in the TNF- α and NF- κ B1 genes with the occurrence of fatigue and sleep disorder in the Parkinson's disease (PD) patients.

2 SUBJECTS AND METHODS

Study Population

A total of 180 PD patients from northeastern Brazil were enrolled in this study. The patients were attended at the PRO-PARKINSON service of the Clinical Hospital of the Federal University of Pernambuco in the period from January 2016 to December 2017. The diagnosis of idiopathic PD was based on the United Kingdom Parkinson's Disease Society Brain Bank criteria [15] and reviewed by an experienced neurologist.

Relevant clinical and demographic features of the study population, such as, sex, age of onset, duration of disease, therapy used, severity of the disease (based on their Hoehn-Yahr score), use of tobacco, alcohol and contact with pesticides, were obtained through medical records and in a face-to-face interview with the patients and their family members. This study was previously evaluated and approved by the ethics committee of the Health Ministry of Brazil (CAAE: 45614415.0.0000.5208) and patients were enrolled after having read and signed an informed consent form.

The occurrence of fatigue was detected when rates were reported for different cut-off points (e.g., ≥ 3.3 and ≥ 8 points on the Parkinson Fatigue Scale [PFS])¹⁶ or was reported according to several recommended measurement scales (e.g., PFS and Fatigue Severity Scale [FSS]) [17]. The sleep disorder score was performed as described by Ray Chaudhuri K et al. [18]. It is a visual scale that addresses 15 symptoms associated with sleep disorder. The maximum cumulative score is 150 (the patient is free of all symptoms).

DNA extraction and genotyping

Genomic DNA was extracted from 3 mL of venous peripheral blood of PD individuals using the Wizard Genomic DNA Purification Kit (Promega, Madison, Wisconsin) and stored in a -20°C freezer. This procedure has been performed in the "Laboratório de Genética Bioquímica e Sequenciamento de DNA Profa. Tânia Falcão (GENOMA)"-Universidade Federal Rural de Pernambuco (UFRPE).

NFKB1 (rs28362491) and TNF- α (rs1800629) variants were genotyped using polymerase chain reaction restriction fragment length polymorphism (PCR-RFLP) assay. A fragment of 289 pb from NFKB1

rs28362491 SNP was amplified using the following set of primers: sense 5'-TTTAATCTGTGAAGAGATGTGAATG-3' and antisense 5' CTCTGGCTTC-CTAGCAGGG and digested by the restriction enzyme *Van91I*, which yields three DNA fragments of 206, 48 and 35 bp for NFKB W allele, two DNA fragments of 254, 35 bp for NFKB D allele [19]. For TNF- α an amplicon of 107 bp was amplified using the following primers: sense 5'-AGGCAATAGGTTTTGAGGGCCAT-3' and antisense 5'- TCCTCCCTGCTGC-GATTCCG-3', that was then digested by restriction enzyme *NcoI* (Fermentas). The GG genotype resulted in two fragments of 87 and 20 bp, while the AA genotype produced one fragment of 107 bp. The GA genotype showed 3 fragments: 107, 87 and 20pb [20]. The digestion products were separated on a 3% agarose gel using the red gel dye and subsequently analyzed on the transilluminator under UV light.

3 STATISTICAL ANALYSES

The genotype distribution and allele frequencies of the polymorphisms were obtained by direct counting. The Hardy-Weinberg equilibrium (HWE) test was applied to datasets using the Bioestat 5.0 program. Univariate and multivariate statistical analyses were performed using the R software, version 3.0.2 (<http://www.R-project.org/>). The association for categorical variables was verified with the chi-squared test. The Student's t-test was performed for continuous data with normal distribution and the Wilcoxon–Mann–Whitney test was performed for data without a normal distribution. The Kolmogorov–Smirnov test was used to evaluate normality. Genotype frequencies were analyzed by the χ^2 test and the odds ratio (OR) with 95% confidence interval (CI) was calculated, using the Bioestat 5.0 program. Data are shown as mean \pm standard deviation (SD) and as median with 25–75th percentiles, as appropriate.

Multiple Poisson regression analyses with robust standard errors were used to assess the effect of polymorphisms on the occurrence of visual hallucinations, since the outcome was a frequent event in our sample [21], while potential confounders to be considered in models were defined based on conceptual analyses of the literature and/or by means of a statistical definition. A formal Bonferroni correction for the number of analyzed SNPs would require a significance threshold of $P = 0.025$.

4 RESULTS

The clinical and demographic characteristics of the patients according to the presence/absence of fatigue and sleep disorder are described in Table 1. In this study, we verified that the prevalence of fatigue and sleep disorder were present in 26.2% and 15% of the patients, respectively. Regarding sleep disorder, there was a significant association with the use of the pramipexole ($p = 0.020$).

Our results of univariate analyses between TNF- α and NFKB polymorphisms (rs1800629 and rs28362491, respectively) and the occurrence of fatigue and sleep disorder are shown in Table 2. No statistical association was found between the genetic variants investigated and the occurrence these side effects ($p > 0.05$).

In contrast, the multivariate analysis revealed an increased risk for the occurrence of fatigue in individuals carrying as the NFKB D/W genotype (Prevalence Ratio: PR=2.091; 95%CI, 1.155-3.784; p=0.015) and D/D genotype (PR= 2.237; 95%CI, 1.078-4.643; p=0.031) (Table 3). In addition, there was an increased risk to the occurrence of fatigue in PD patients with sleep disorder (PR= 5.226; 95%CI, 2.934-9.307; p= 0.000). Regarding the TNF- α gene, no significant association was observed between the occurrence of fatigue and/or sleep disorder (p > 0.05) (Table 3).

5 DISCUSSION

Although studies had showed association of inflammation with Parkinson's disease, the mechanisms where by it contributes to the progression of the disease are still unclear [22]. Experimental and postmortem studies suggest that chronic inflammation associated with microglial activation could act as a stressor event promoting the recruitment of pro-inflammatory molecules such as tumor necrosis factor (TNF α) and nuclear factor kappa B (NFKB1) contributing to the progressive degeneration of dopaminergic neurons [23] and the appearance of non-motor symptoms such as fatigue and sleep disorders, once this symptoms may be due alterations in inflammatory molecules.

Fatigue in neurological diseases has been described as a significant decrease in energy level ou an increased persecution of efforts disproportionate to attempted activities [24]. Friedman et al. (2018) reported that on-third of their patients rated fatigue as one of the most disabling symptoms of Parkinson's disease and more than 50% of the patients assessed had this symptom [25]. Others studies in different populations, however, reported prevalence ranging from 33% to 58% of the patients studied [2, 23, 24-31]. In our study the prevalence of fatigue was 26.2% suggesting that this difference can be explained once fatigue is assessed as categorical variable (presence or absence).

In general, fatigue is defined as a feeling of exhaustion for a certain period of time and is not explained by the effects of drugs, medical or psychiatric disorders [24]. In Parkinson's disease, fatigue is considered one of the most common and incapacitating non-motor symptoms, and may manifest itself even in the pre-motor stages of the disease and, once present, may persist and/or worsen over time [26,27].

Previous reports showed that the prevalence of sleep disorder ranged 50% to 81% [29,32–34]. In the present study, only 15% of the patients evaluated presented sleep disturbance. Again, we can suggest that this difference can be explained once sleep disorder is assessed as categorical variable (presence or absence).

Sleep disorder was described by James Parkinson as severe exhaustion and extreme drowsiness, and although the onset of motor symptoms in the PD often preceding non-motor symptoms, some patients may begin to exhibit non-motor symptoms in the course of the disease [2].

According to Titova and colleagues (2017) in a study evaluating non-motor characteristics of Parkinson's disease, the main treatments for non-motor disease symptoms are from non-pharmacological origin [35]. However, motor symptoms usually precede the non-motor symptoms of PD and have as main

treatment the use of drugs that, in the long time, may lead to the onset of adverse events, as well as triggering new symptoms of the disease [2,36–38]. In this line of reasoning, our results showed association between the use of pramipexole and the development of sleep disturbance ($p = 0.020$).

The transcription factor NF κ B1 was first discovered on T lymphocytes [39], but is present in all mammalian cells. The most well-known and studied role of this transcription factor is in the immune response, in which it regulates expression of essential genes in the inflammatory process and defense against parasites. In addition, it also acts on cell survival and proliferation, apoptosis and CNS function [40]. The TNF α protein initially known for its cytotoxicity, was originally identified in mouse serum after injection with *Mycobacterium bovis* Calmette-Guerin (BCG) bacillus and endotoxin. It is a proinflammatory cytokine, encoded by the TNF- α gene, that actively regulates a broad spectrum of biological processes, notably neuroinflammation [41,42].

NF κ B1 is a transcription factor that modulates the balance of pro and anti-inflammatory cytokines. Wang et al. showed that the polymorphism -94 ins / del ATTG (rs28362491) involving the deletion of four nucleotides in the promoter region of the NF κ B1 gene consists of three genotypes: homozygous or wild type (WW), homozygous or variant (DD); heterozygous ins / del (DW) and is within the promoter region of the NF κ B1 gene, potentially affecting gene transcription and NF κ B protein function, leading sequentially to loss of nuclear protein binding capacity and reduced activity.

Individuals carrying the mutant genotype may have the activity of this gene reduced and thus not be able to respond adequately to certain stimuli in order to aid in the transcription of genes whose products would act in an attempt to control the inflammatory process, and may end up making it chronic and aiding in the onset of various physiological complications [43]. These information can explain our data from multivariate analysis that revealed that individuals carrying DW or DD genotypes for NF κ B gene had 2-fold increased risk for the occurrence of fatigue [31].

Recently, Herlofson et al. [31] showed that the development of fatigue would be associated with changes in metabolism, endocrine system and inflammatory markers and it is not related to the central nervous system. These information can explain our data from multivariate analysis that revealed that individuals carrying DW or DD genotypes for NF κ B gene had 2-fold increased risk for the occurrence of fatigue.

Despite of Lindqvist et al. [44] had found significant and positive correlation between high level of TNF α and the presence of non-motor symptoms such as depression, anxiety and fatigue, no significant association between TNF α polymorphism and occurrence of fatigue was observed in the present study. Our data suggest that this polymorphism could not be is associated with the occurrence of fatigue. To the best of our knowledge this is the first study that correlated fatigue and sleep disorder in Parkinson's disease with polymorphisms in the promoter region of TNF α and NF κ B1 genes (rs1800629 and rs28362491, respectively).

6 CONCLUSION

In conclusion, our study suggested that NFKB1 variant and use of the pramipexole may have significant influence on increasing risk for the occurrence of fatigue and sleep disorder, respectively in a Brazilian PD population.

ACKNOWLEDGEMENT

We thank the patients and their families, whose collaboration and understanding have made this work possible.

Funding

Our study was supported by the Brazilian funding agency FACEPE (Fundação de Amparo à Ciência e Tecnologia do Estado de Pernambuco).

Conflict of interest

The authors declare that they have no competing interests.

Consent to participate

The work was done according to all ethical principles and criteria of good scientific practice and data used from official databases that were processed statistically. The study does not contain personal data about patients nor does it include animal studies.

Consent to publish

All authors agree with the publication of the research.

REFERENCES

1. Bassani TB et al. Neuroinflammation in the pathophysiology of Parkinson's disease and therapeutic evidence of anti-inflammatory drugs. *Arq Neuropsiquiatr* 2015; 73(7): 616–623. doi:10.1590/0004-282X20150057.
2. Kalia LV, Lang AE. Parkinson's disease. *The Lancet* 2015; 386(9996): 896–912. doi:10.1016/S0140-6736(14)61393-3.
3. Klein C, Schlossmacher MG. Parkinson disease, 10 years after its genetic revolution: Multiple clues to a complex disorder. *Neurology* 2007; 69(22): 2093–2104. doi:10.1212/01.wnl.0000271880.27321.a7.
4. Hirsch EC et al. Neuroinflammation in Parkinson's disease. *Parkinsonism Relat Disord* 2012; 18: S210–S212. doi:10.1016/S1353-8020(11)70065-7.
5. Ghosh A et al. Selective inhibition of NF- κ B activation prevents dopaminergic neuronal loss in a mouse model of Parkinson's disease. *Proc Natl Acad Sci* 2007; 104(47): 18754–18759. doi:10.1073/pnas.0704908104.
6. Agliardi C et al. TNF- α – 308 G/A and – 238 G/A promoter polymorphisms and sporadic Parkinson's disease in an Italian cohort. *J Neurol Sci* 2018; 385: 45–48. doi:10.1016/j.jns.2017.12.011.
7. Carvey PM et al. Intra-parenchymal injection of tumor necrosis factor- α and interleukin 1- β produces dopamine neuron loss in the rat. *J Neural Transm* 2005; 112(5): 601–612. doi:10.1007/s00702-004-0222-z.
8. Nagatsu T, Sawada M. Inflammatory process in Parkinson's disease: role for cytokines. *Curr Pharm Des* 2005; 11(8): 999–1016. doi: 10.2174/1381612053381620.
9. Dantzer R et al. From inflammation to sickness and depression: when the immune system subjugates the brain. *Nat Rev Neurosci* 2008; 9(1): 46–56. doi:10.1038/nrn2297.
10. Irwin MR, Cole SW. Reciprocal regulation of the neural and innate immune systems. *Nat Rev Immunol* 2011; 11(9): 625–632. doi:10.1038/nri3042.
11. Miller AH et al. Neuroendocrine-immune mechanisms of behavioral comorbidities in patients with cancer. *J Clin Oncol Off J Am Soc Clin Oncol* 2008; 26(6): 971–982. doi:10.1200/JCO.2007.10.7805.
12. Ammal Kaidery N et al. Epigenetic Landscape of Parkinson's Disease: Emerging Role in Disease Mechanisms and Therapeutic Modalities. *Neurotherapeutics* 2013; 10(4): 698–708. doi:10.1007/s13311-013-0211-8.
13. Gius D et al. Distinct effects on gene expression of chemical and genetic manipulation of the cancer epigenome revealed by a multimodality approach. *Cancer Cell* 2004; 6(4): 361–371. doi:10.1016/j.ccr.2004.08.029.
14. Chen M, Zhang L. Epigenetic mechanisms in developmental programming of adult disease. *Drug Discov Today* 2011; 16(23–24): 1007–1018. doi:10.1016/j.drudis.2011.09.008.
15. Hughes AJ et al. What features improve the accuracy of clinical diagnosis in Parkinson's disease: a clinicopathologic study. *Neurology* 1992; 42(6): 1142–1146.
16. Kummer A, Scalzo P, Cardoso F, Teixeira AL. Evaluation of fatigue in Parkinson's disease using the Brazilian version of Parkinson's Fatigue Scale. *Acta Neurol Scand.* 2011 Feb;123(2):130-6. doi: 10.1111/j.1600-0404.2010.01364.x.
17. Solla P et al. Association between fatigue and other motor and non-motor symptoms in Parkinson's disease patients. *J Neurol* 2014; 261(2): 382–391. doi:10.1007/s00415-013-7207-5.

18. Chaudhuri KR et al. The Parkinson's disease sleep scale: a new instrument for assessing sleep and nocturnal disability in Parkinson's disease. *J Neurol Neurosurg Psychiatry* 2002; 73(6): 629–635. doi: 10.1136/jnnp.73.6.629.
19. Salim PH et al. Interleukin-10 Gene Promoter and NFKB1 Promoter Insertion/Deletion Polymorphisms in Systemic Sclerosis. *Scand J Immunol* 2013; 77(2): 162–168. doi:10.1111/sji.12020.
20. Tavares MC, de Lima Júnior SF, Coelho AV, Marques TR, de Araújo DH, Heráclio Sde A, Amorim MM, de Souza PR, Crovella S. Tumor necrosis factor (TNF) alpha and interleukin (IL) 18 genes polymorphisms are correlated with susceptibility to HPV infection in patients with and without cervical intraepithelial lesion. *Ann Hum Biol.* 2016 May;43(3):261-8. doi: 10.3109/03014460.2014.1001436.
21. Greenland S. Model-based Estimation of Relative Risks and Other Epidemiologic Measures in Studies of Common Outcomes and in Case-Control Studies. *Am J Epidemiol* 2004; 160(4): 301–305. doi:10.1093/aje/kwh221.
22. Jang H et al. Viral parkinsonism. *Biochim Biophys Acta BBA - Mol Basis Dis* 2009; 1792(7): 714–721. doi:10.1016/j.bbadis.2008.08.001.
23. Tansey MG, Goldberg MS. Neuroinflammation in Parkinson's disease: Its role in neuronal death and implications for therapeutic intervention. *Neurobiol Dis* 2010; 37(3): 510–518. doi:10.1016/j.nbd.2009.11.004.
24. Kluger BM et al. Fatigue and fatigability in neurologic illnesses: Proposal for a unified taxonomy. *Neurology* 2013; 80(4): 409–416. doi:10.1212/WNL.0b013e31827f07be.
25. Friedman JH et al. Fatigue in Parkinson's disease: report from a multidisciplinary symposium. *Npj Park Dis* 2016; 2(1). doi:10.1038/npjparkd.2015.25.
26. Barone P et al. The PRIAMO study: A multicenter assessment of nonmotor symptoms and their impact on quality of life in Parkinson's disease. *Mov Disord* 2009; 24(11): 1641–1649. doi:10.1002/mds.22643.
27. Pont-Sunyer C et al. The Onset of Nonmotor Symptoms in Parkinson's disease (The ONSET PDStudy): The Onset PD Study. *Mov Disord* 2015; 30(2): 229–237. doi:10.1002/mds.26077.
28. Siciliano M et al. Fatigue in Parkinson's disease: A systematic review and meta-analysis: Fatigue in Parkinson's Disease. *Mov Disord* 2018; 33(11): 1712–1723. doi:10.1002/mds.27461.
29. van Hilten JJ et al. Diurnal effects of motor activity and fatigue in Parkinson's disease. *J Neurol Neurosurg Psychiatry* 1993; 56(8): 874–877.
30. Stocchi F et al. Prevalence of fatigue in Parkinson disease and its clinical correlates. *Neurology* 2014; 83(3): 215–220. doi:10.1212/WNL.0000000000000587.
31. Herlofson K, Kluger BM. Fatigue in Parkinson's disease. *J Neurol Sci* 2017; 374: 38–41. doi:10.1016/j.jns.2016.12.061.
32. Bollu PC, Sahota P. Sleep and Parkinson Disease. *Mo Med* 2017; 114(5): 381–386.
33. Nausieda PA et al. Sleep disruption in the course of chronic levodopa therapy: an early feature of the levodopa psychosis. *Clin Neuropharmacol* 1982; 5(2): 183–194. doi: 10.1097/00002826-198205020-00003.
34. Riedel O et al. Frequency of dementia, depression, and other neuropsychiatric symptoms in 1,449 outpatients with Parkinson's disease. *J Neurol* 2010; 257(7): 1073–1082. doi:10.1007/s00415-010-5465-z.
35. Titova N, Chaudhuri KR. Personalized medicine in Parkinson's disease: Time to be precise: Parkinson's and Personalized Medicine. *Mov Disord* 2017; 32(8): 1147–1154. doi:10.1002/mds.27027.

36. Van Gerpen JA et al. Levodopa-Associated Dyskinesia Risk Among Parkinson Disease Patients in Olmsted County, Minnesota, 1976-1990. *Arch Neurol* 2006; 63(2): 205. doi:10.1001/archneur.63.2.205.
37. Fox SH, Lang AE. Levodopa-related motor complications--phenomenology. *Mov Disord Off J Mov Disord Soc* 2008; 23 Suppl 3: S509-514. doi:10.1002/mds.22021.
38. Ku S, Glass GA. Age of Parkinson's disease onset as a predictor for the development of dyskinesia. *Mov Disord* 2010; 25(9): 1177–1182. doi:10.1002/mds.23068.
39. Nabel G, Baltimore D. An inducible transcription factor activates expression of human immunodeficiency virus in T cells. *Nature* 1987; 326(6114): 711–713. doi:10.1038/326711a0.
40. Xiao W. Advances in NF-kappaB signaling transduction and transcription. *Cell Mol Immunol* 2004; 1(6): 425–435.
41. Shirai T et al. Cloning and expression in *Escherichia coli* of the gene for human tumour necrosis factor. *Nature* 1985; 313(6005): 803–806. doi: 10.1038/313803a0.
42. Pennica D et al. Human tumour necrosis factor: precursor structure, expression and homology to lymphotoxin. *Nature* 1984; 312(5996): 724–729. doi: 10.1038/312724a0.
43. Wang D et al. Genetic association between NFKB1 -94 ins/del ATTG Promoter Polymorphism and cancer risk: a meta-analysis of 42 case-control studies. *Sci Rep* 2016; 6: 30220. doi:10.1038/srep30220.
44. Lindqvist D et al. Non-Motor Symptoms in Patients with Parkinson's Disease – Correlations with Inflammatory Cytokines in Serum. Duda J, ed. *PLoS ONE* 2012; 7(10): e47387. doi:10.1371/journal.pone.0047387.

Table 1. Clinical and demographic characteristics of Brazilian PD patients: all and stratified for fatigue and sleep disorder.

	All	Fatigue		P (value)	Sleep Disorder		P (value)
		No	Yes		No	Yes	
Patients (%)	180	133 (73.8)	47 (26.2)		153 (85%)	27 (15%)	
Men (%)	107 (59.4)	79 (59.4)	28 (40.6)	0.879	91 (59.5)	16 (40.5)	0.848
Age at onset	57.5	59	54	0.129	58	53	0.108
Median (range) years	(48-64)	(48-66)	(49-60)		(48-65)	(48-61)	
Disease Duration	7.68±	7	7	0.979	7	7	0.312
Mean± s.d.	4.06	(5-10)	(5-9.5)		(5-9)	(5.5-12)	
Levodopa Therapy duration	6.72±	6	6	0.880	6	6	0.269
Mean ± s.d.	3.74	(4-9)	(4-8)		(4-9)	(4-9.5)	
Levodopa dosage	720.13±	600	800	0.208	600	600	0.961
Mean ±s.d.	312.16	(500-800)	(600-1000)		(600-900)	(500-900)	
HY stage (%)							
1(%)	23(12.8)	19(14.3)	4(8.5)	0.444	21(13.7)	2(7.5)	0.552
2(%)	91(50.5)	67(50.4)	24(51.0)	0.929	80(52.3)	11(40.7)	0.369
3(%)	45(25)	29(21.7)	16(34.0)	0.141	34(22.2)	11(40.7)	0.070
4(%)	18(10)	15(11.3)	3(6.5)	0.497	15(9.8)	3(11.1)	0.889
5(%)	3(1.7)	3(2.3)	0(0.0)	0.707	3(2.0)	0(0.0)	0.935
Pramipexole	108(60.0)	79(59.4)	29(61.7)	0.917	87(56.9)	21(77.7)	0.020
Smoking	20(11.1)	14(10.5)	6(12.7)	0.880	18(11.8)	2(7.4)	0.739
Alcoholism	22(12.2)	15(11.3)	7(14.9)	0.695	19(12.4)	3(11.1)	0.898
Agrotoxic	22(12.2)	13(9.7)	9(19.1)	0.153	18(11.8)	4(14.8)	0.898

± SD, standard derivation. P (values) calculated by t test or by Wilcoxon–Mann–Whitney U-test (quantitative variables with or without normal distribution, respectively), and X^2 test (categorical variables). Significant p-values are shown in bold.

Table 2. Univariate analyses showing the relation between *TNF α* rs1800629, *NFKB1* rs28362491 and the occurrence of fatigue and sleep disorder in Brazilian PD patients.

	Total	Fatigue		X ² (p)	OR	P*	Sleep Disorder		X ² (p)	OR (IC 95%)	P*		
		No (133)	Yes (47)				No (153)	Yes (27)					
<i>TNFα</i>													
G/G	119(66.0)	87 (65.4)	32 (68.1)	3.25 (0.20)	1.00	0.21	101(66.0)	18 (67.0)	0.83 (0.66)	1.00	0.62		
G/A	47(26.0)	38 (28.6)	9 (19.1)				0.64(0.28-1.48)	39 (25.0)				8 (30.0)	1.15 (0.46-2.86)
A/A	14(8.0)	8 (6.0)	6 (12.8)				2.04(0.66-6.33)	13 (8.0)				1 (4.0)	0.43 (0.05-3.51)
<i>NFKB1</i>													
W/W	48(26.7)	39 (29.3)	9 (19.1)	2.31 (0.32)	1.00	0.31	38 (25.0)	10 (37.0)	2.33 (0.31)	1.00	0.32		
W/D	89(49.5)	65 (48.9)	24 (51.1)				1.60 (0.68-3.79)	79 (52.0)				10 (37.0)	0.48 (0.18-1.25)
D/D	43(23.8)	29 (21.8)	14 (29.8)				2.09 (0.80-5.49)	36 (24.0)				7 (26.0)	0.74 (0.26-2.15)

Abbreviations: P*: P values; X²: Chi- square.

Table 3. Multiple Poisson regression model adjusted for clinical variables and *TNFA* (rs1800629) and *NFKB1* (rs28362491) polymorphisms predicting fatigue in Parkinson's disease patients.

	Prevalence Ratio	95% CI	p - value
Gender (Male)	0.790	0.479-1.304	0.357
rs1800629		0.90-1.46	0.241
A/A	2.083	0.920-4.716	0.078
G/A	0.646	0.358-1.166	0.147
rs 28362491			
D/D	2.237	1.078-4.643	0.031
D/W	2.091	1.155-3.784	0.015
Age	0.991	0.966-1.016	0.474
Disease duration (years)	0.951	0.787-1.151	0.608
Levodopa therapy duration	1.026	0.857-1.229	0.778
PDSS	5.226	2.934-9.307	0.000
HY	0.974	0.651-1.456	0.897
Levodopa dose	1.001	1.0-1.001	0.131
Dopamine Agonists			
Pramipexole use	0.793	0.483-1.300	0.357
Pain	0.624	0.149-2.617	0.519
Bradykinesia	1.147	0.400-3.290	0.798
Rigidity	1.020	0.501-2.075	0.957
Tremor	1.335	0.563-3.167	0.512
Smoking	1.063	0.485-2.328	0.878
Alcoholism	1.319	0.660-2.640	0.433
Agrotoxic	1.581	0.979-2.554	0.061

Abbreviations: .PDSS: PD Sleep scale; HY: Hoehn and Yahr. Significant p-values are shown in bold.