



<https://doi.org/10.56238/emerrelcovid19-064>

Caroline Brum Gonzalez

Master's Degree, Federal University of Rio de Janeiro/UFRJ

Senda Charone

PhD, University of São Paulo/USP

Elizeu Fagundes de Carvalho

Post Doctorate, Washington University - WashU

Sonia Groisman

Post Doctorate, American World University, USA

1 INTRODUCTION

In December 2019, the global spread of a new virus known as SARS-CoV-2, which is responsible for the disease called COVID-19, occurred, triggering an unprecedented crisis around the world. On March 11, 2020, the World Health Organization (WHO) declared the situation a pandemic. As of June 16, 2020, there have been more than 7.9 million confirmed cases and 434,796 deaths in more than 200 countries. Brazil was the second most affected country, with more than 900,000 confirmed cases and 43,332 deaths.

As a measure to contain the rapid spread of COVID-19, isolation and social distancing measures have been implemented around the world. This resulted in the temporary closure of dental clinics, both public and private, and the suspension of elective dental care. In addition, the dental school clinics were closed due to the interruption of face-to-face activities and clinical teaching.

Considering the current knowledge about the biological behavior of COVID-19, it is important to note that many viral infections can cause oral manifestations. Therefore, it is valid to know the oral manifestations that may be a direct consequence of SARS-CoV-2 infection.

There are several oral manifestations related to COVID-19, which can occur due to the side effects of the medications used in the treatment of the disease, as well as conditions arising from mechanical ventilation therapy and hospitalization in intensive care units (ICU). These manifestations include changes in the characteristics of the mucous membranes, changes in the production and quality of saliva, stomatitis, ulcers, sensory changes, pigmentation, among others. The advanced age and severity of COVID-19 can aggravate these injuries. Other factors, such as lack or reduction of oral hygiene, stress, systemic diseases (such as diabetes mellitus and immunosuppression), trauma from intubation, vascular impairment, and hyperinflammatory response secondary to COVID-19, may also contribute to the development of these lesions.

2 ORAL MANIFESTATIONS ASSOCIATED WITH COVID-19

2.1 CHANGES IN TASTE

Dysgeusia is a persistent distortion of taste, whether transient or permanent. Dysgeusia is also often associated with ageusia, which is the complete lack of taste, and hypogeusia, which is a decreased sensitivity to taste. The oral symptoms of COVID-19 are characterized by disturbances in taste and smell, without the simultaneous presence of nasal congestion. SARS-CoV-2 has been found to bind to the epithelial cells of the tongue via the converting enzyme receptor, which is widely expressed in these cells. Based on this finding, dysgeusia is considered the first direct oral manifestation of the infection.

2.2 HYPOSALIVATION

Additionally, decreased saliva production (hyposalivation) or changes in saliva composition can lead to dysgeusia and even ageusia. A recent study demonstrated that more than 50% of COVID-19 patients had symptoms of dry mouth (xerostomia) and dysgeusia, and there was a significant correlation between these two symptoms. However, more research is needed to determine if taste dysfunction is a direct consequence of SARS-CoV-2 infection or if the virus causes hyposalivation, resulting in dysgeusia. Either way, one symptom aggravates the other. Reducing saliva production also increases the risk of caries injuries and aggravating the process of tooth decay. In addition, the decrease in saliva implies a lower amount of immunoglobulins, which can pose a great risk for the development of periodontal diseases

2.3 OPPORTUNISTIC INFECTIONS

It is essential to highlight that the acute condition caused by COVID-19 can increase susceptibility to opportunistic infections, such as herpes simplex and candidiasis, which may present similar clinical characteristics. Since there is no standard treatment for COVID-19, different medications are given to patients, and it is important to consider that other endemic infectious diseases may arise as a consequence of COVID-19.

Studies indicate that prolonged use of antibiotics may aggravate oral manifestations and warn of the risk of potentially fatal opportunistic oral infections resulting from the indiscriminate prescription of broad-spectrum antibiotics. Mucus skin complications in the oral cavity can arise due to several factors, such as prolonged positioning of mechanical ventilation devices in intensive care units (ICU), previous antibiotic treatments that can lead to immune dysregulation, and, in general, injuries are indicative of side effects related to deteriorating systemic health or the use of medications in the treatment of COVID-19.



Source: Internet - Opportunistic Infection: Herpes Simplex



Source: Internet - Opportunistic Infection: Oral Candidiasis

2.4 MUCOSAL LESIONS

Several isolated case reports or case series have documented oral manifestations of COVID-19. However, questions arise as to whether these lesions are a direct consequence of SARS-CoV-2 infection or whether they are secondary manifestations of the disease. These manifestations include oral ulcers, petechiae, and reddish macules, mainly on the tongue and palate.

The discussion about the origin is inconclusive, since it may be the virus as to the pharmacological treatment employed. According to recent studies, the most frequent oral lesions were aphthous ulcerations (65%), vesicular-bullous lesions (11%), macules (16%), petechiae (6%) and plaques (6%), most of which were symptomatic, located on the tongue (53%), palate (30%), lip (23%) and gums (5%). Other oral changes reported were ageusia/dysgeusia (21%), burning in the mouth (20%) and xerostomia/hyposalivation (7%).



Source: Internet - Mucosal Lesions: Canker sores



Source: Internet - Mucosal Lesions: Petechiae

3 CONCLUSION

It cannot be stated with certainty that the oral manifestations are directly related to the SARS-CoV-2 virus or if they are side effects of the treatment of the disease. While it is common for COVID-19 patients to experience oral signs and symptoms, such as traumatic ulcers, changes in taste, opportunistic infections, and reduced salivary flow, these manifestations are not mandatory. Understanding the mechanism of action of COVID-19 requires more scientific research to determine the origin of these lesions and oral manifestations, as well as find ways to minimize them. It is important to make dentists aware of the proper management of the patient in this context.

REFERENCES

Organização mundial da saúde. Painel da oms sobre a doença por coronavírus (covid-19). Disponível em: <https://who.sprinklr.com> [acesso em 16 de junho de 2020]. » <https://who.sprinklr.com>

Perez dec, passos kkm, machado ra, martelli-junior h, bonan prf. Educação continuada em câncer bucal durante surto de coronavírus 2019 (covid-19). *Oral oncol* 2020; 105:104713. <https://doi.org/10.1016/j.oraloncology.2020.104713>
» <https://doi.org/10.1016/j.oraloncology.2020.104713>

Chary e, carsuzaa f, trijolet jp, capitaine al, roncato-sabera m, fouet k, et al. *Sou j rhinol alerg* 2020; 1945892420930954. <https://doi.org/10.1177/1945892420930954>
» <https://doi.org/10.1177/1945892420930954>

Xu h, zhong l, deng j, peng j, dan h, zeng x, et al. *Int j oral sci* 2020; 12(1):8. <https://doi.org/10.1038/s41368-020-0074-x>» <https://doi.org/10.1038/s41368-020-0074-x>

Mariz bala, marcatb marca, ribeiro acp, lopes ma, santos-silva ar. *Novos insights para a patogênese da disgeusia relacionada à covid-19*. *j dent res* 2020; 2020:22034520936638. <https://doi.org/10.1177/0022034520936638>» <https://doi.org/10.1177/0022034520936638>

Biadsee a, biadsee a, kassem f, dagan o, masarwa s, ormianer z. *Manifestações olfativas e orais de covid-19: sintomas relacionados ao sexo - um caminho potencial para o diagnóstico precoce*. *Otorrinolaringologia cirurgia de cabeça e pescoço* 2020; 194599820934380 <https://doi.org/10.1007/s00405-020-05965-1>

Pitts nb, zero dt, ekstram k, weintraub já, ramos-gomesf, tagami j, twetman s, tsakos g, ismail a. *dental caries*. *nat rev dis primers*. 2007;3:17030.

Soares cd, carvalho ra, carvalho ka, carvalho mg, almeida op. *Lesões bucais em paciente com covid-19*. *Med oral patol oral cir bucal* 2020; 2020:240-44. <https://doi.org/10.4317/medoral.24044>» <https://doi.org/10.4317/medoral.24044>

Santos jad, normando agc, silva rlcd, paula rm, cembranel ac, santos-silva ar, et al. *Lesões da mucosa oral em paciente com covid-19: novos sinais ou manifestações secundárias?* *Int j infect dis* 2020; 97:326-8. <https://doi.org/10.1016/j.ijid.2020.06.012>
» <https://doi.org/10.1016/j.ijid.2020.06.012>

Rocha ba, soto gr, grossmann smc, aguiar mcf, andrade bab, romañach mj, horta mcr. *Enantema viral na mucosa oral: um possível desafio diagnóstico na pandemia de covid-19*. *Oral dis* 2020; 10.1111/13473. <https://doi.org/10.1111/odi.13473>» <https://doi.org/10.1111/odi.13473>
» <https://doi.org/10.1111/odi.13473>

Chiotos k, bassiri h, behrens em, blatz am, chang j, diorio c, et al. *J pediátrica infect dis soc*. 2020; p1aa069. <https://doi.org/10.1093/jpids/p1aa069>
» <https://doi.org/10.1093/jpids/p1aa069>

Clothier b, stringer m, jeffcoat mk. *Doença periodontal e desfechos da gravidez: exposição, risco e intervenção*. *Melhor pract res clin obstet gynaecol*. 2007;21(3):451-466.

Cuidados com a saúde bucal durante a gestação e ao longo da vida. *Obsteto ginecologista*. 2013;122(2 pt 1):417-422.

M maltz, lma tenuta, s groisman, ja cury . *Cariologia: conceitos básicos, diagnóstico e tratamento não restaurador: série abeno: odontologia essencial-parte clínica*. 1st edition, artes medicas, 2016, 143 pags.

http://www.gcamerica.com/products/preventive/saliva_check_buffer/index.php

Marsh pd. Placa dentária como biofilme e comunidade microbiana: implicações para a saúde e a doença. *Bmc saúde bucal* 2006; 6(suppl. 1): s14. [<http://dx.doi.org/10.1186/1472-6831-6-s1-s14>] [pmid: 16934115]

Kolenbrander pe, andersen rn. Caracterização da coagregação mediada por adesina *Streptococcus gordonii* (s. Sanguis) pk488 com *Actinomyces naeslundii* pk606. *Infect immun* 1990; 58(9): 3064-72. [pmid: 2387635]

Marsh pd. Ecologia microbiana da placa dental e sua importância na saúde e na doença. *Adv dent res* 1994; 8(2): 263-71. [<http://dx.doi.org/10.1177/08959374940080022001>] [pmid: 7865085]

Lynge pedersen, a.m.; belstrom, d.; o papel das defesas salivares naturais na manutenção de uma microbiota oral saudável. *J. Dent*. 2019, 80 (suppl. 1), s3–s12. [crossref]

Siddiqui, h.; oliveira, a.; oliveira, a.; pereira, m.m.; oliveira, r.; olsen, i. Análise microbiológica e bioinformática de pacientes com síndrome de sjögren primária com salivagem normal. *J. Microbiol oral*. 2016, 8, 31119. [crossref]

De paiva, c.s.; pereira, d.b.; ribeiro, m.e.; ribeiro, f.; pereira, j.l.; oliveira, a.; ribeiro, c.f.; oliveira, d.s.; oliveira, j.n.; oliveira, j.f.; diversidade alterada do microbioma mucoso e gravidade da doença na síndrome de sjögren. *Rep*. 2016, 6, 23561. [crossref]

Zhou, z.; oliveira, g.; ribeiro, a.; ribeiro, z.; oliveira, c.; oliveira, a.; chen, x. Análise molecular da microflora oral em pacientes com síndrome de sjögren primária usando sequenciamento de alto rendimento. *Peerj* 2018, 6, e5649. [crossref]

Van der meulen, t.a.; oliveira, j.j.m.; ribeiro, a.; oliveira, a.c.; oliveira, a.; oliveira, a.; pereira, j.; pereira, c.; ribeiro, f.k.l.; oliveira, f.g.m.; et al. Disbiose do microbioma da mucosa bucal em pacientes com síndrome de sjögren primária. *Reumatologia (oxford)* 2018, 57, 2225–2234. [crossref]

Leung, k.c.; oliveira, j.j.; mcmillan, a.s. microbiota supragengival na síndrome de sjögren. *Clin. Investigação oral*. 2007, 11, 415–423. [crossref]

Rodrigues, s.; oliveira, a.k.; oliveira, a.; oliveira, a.g.; oliveira, e.b.; oliveira, o.; oliveira, m.; jensen, j.l. microbiota salivar disbiótica em pacientes com boca seca e síndrome de sjögren primária. *Plos one* 2019, 14, e0218319. [crossref]

Szymula, a.; oliveira, j.; oliveira, m.b.; ribeiro, a.; pereira, a.m.; deshmkh, epítipo de células t dos eua mimetismo entre a síndrome de sjögren antígeno a (ssa)/ro60 e bactérias orais, intestinais, cutâneas e vaginais. *Clin. Immunol*. 2014, 152, 1–9. [crossref]

Sudhakara, p.; ribeiro, a.; ribeiro, a.; wilson, a. Comunidades disbióticas orais e suas implicações em doenças sistêmicas. *Dent. J*. 2018, 6, 10. [crossref]

Frank, j. A., reich, c. I., sharma, s., weisbaum, j. S., wilson, b. A., e olsen, g. J. (2008). Avaliação crítica de dois primers comumente utilizados para amplificação de genes bacterianos do 16s rrna. Appl. Meio ambiente. Microbiol. 74, 2461–2470. Doi: 10.1128/aem.02272-07

Lane, d. J., pace, b., olsen, g. J., stahl, d. A., sogin, m. L., e pace, n. R. (1985). Determinação rápida de sequências de rna ribossomal 16s para análises filogenéticas. Proc. 82, 6955–6959. Doi: 10.1073/pnas.82.20.6955

Al-hebshi, n. N., alharbi, f. A., mahri, m., e chen, t. (2017a). Diferenças no bacterioma de produtos de tabaco sem fumaça com diferentes carcinogenicidades orais: análise composicional e funcional predita. Genes (basileia) 8 (4), 106. Doi: 10.20944/preprints201703.0030.v1 .

Al-hebshi, n. N., nasher, a. T., maryoud, m. Y., homeida, h. E., chen, t., idris, a. M., et al. Bacterioma inflamatório com fusobacterium nucleatum e pseudomonas aeruginosa identificado em associação com carcinoma epidermóide oral. Rep. 7, 1834. Doi: 10.1038/s41598-017- 02079-3.

Fronteiras em microbiologia celular e de infecção | www.frontiersin.org 11 novembro 2020 | volume 10 | artigo 602239 frid et al.