

## INCLUSION OF THE GIANT MEALWORM (ZOPHOBAS MORIO) AS AN ALTERNATIVE PROTEIN SOURCE IN DIETS FOR *P. VANNAMEI* SHRIMP (BOONE, 1931)

## INCLUSÃO DO TENÉBRIOS GIGANTE (ZOPHOBAS MORIO) COMO FONTE PROTEICA ALTERNATIVA EM DIETAS PARA CAMARÃO *P. VANNAMEI* (BOONE, 1931)

## INCLUSIÓN DEL GUSANO DE LA HARINA GIGANTE (ZOPHOBAS MORIO) COMO FUENTE ALTERNATIVA DE PROTEÍNA EN DIETAS PARA CAMARONES *P. VANNAMEI* (BOONE, 1931) DE LA APLICACIÓN SHEET2PROMPT API



<https://doi.org/10.56238/sevened2025.036-106>

**Vanuza de Paula do Nascimento da Silva<sup>1</sup>, Fernanda Reis Lima<sup>2</sup>, Léa Carolina de Oliveira Costa<sup>3</sup>, Saymon Rodrigues Matos da Costa<sup>4</sup>, Jackson Oliveira Andrade<sup>5</sup>, Danilo Acatauassu da Silva Costa<sup>6</sup>, Lian Valente Brandão<sup>7</sup>**

### ABSTRACT

This study investigates the inclusion of giant mealworms (*Zophobas morio*) as an alternative protein source in diets for the shrimp *Penaeus vannamei* (Boone, 1931). The main objective is to evaluate the effects of this inclusion on the growth performance of the shrimp. The methodology involved the formulation of experimental diets with different levels of *Zophobas morio* inclusion and the performance of feeding trials under controlled conditions. The results indicated that the inclusion of giant mealworms can completely replace fishmeal without compromising shrimp performance, in addition to presenting additional benefits in terms of health and environmental sustainability. It is concluded that *Zophobas morio* is a viable and promising alternative for formulating shrimp diets, contributing to the diversification of protein sources in aquaculture.

**Keywords:** *Zophobas morio*. Alternative Protein. *Penaeus vannamei*.

### RESUMO

Este trabalho investiga a inclusão do Tenébrio gigante (*Zophobas morio*) como uma fonte proteica alternativa em dietas para o camarão *Penaeus vannamei* (Boone, 1931). O objetivo

<sup>1</sup> Graduated in Fisheries Engineering. Instituto Federal do Pará. E-mail: 2001vpaula@gmail.com  
Lattes: <https://lattes.cnpq.br/2831154724216585>

<sup>2</sup> Graduated in Fisheries Engineering. Instituto Federal do Pará. E-mail: fernandalinma23@gmail.com  
Lattes: <https://lattes.cnpq.br/9068615256466042>

<sup>3</sup> Dr. in Aquaculture. Universidade Federal de Rio Grande (FURG). E-mail: leacarinacosta@yahoo.com.br  
Orcid: <https://orcid.org/0000-0002-4423-7937> Lattes: <https://lattes.cnpq.br/7576540554112066>

<sup>4</sup> Master's degree in Aquaculture and Tropical Aquatic Resources. E-mail: saymon.costa@ifpa.edu.br  
Orcid: 0009-0000-2120-5185 Lattes: <http://lattes.cnpq.br/8426042496733123>

<sup>5</sup> Aquaculture Technologist. Instituto Federal do Pará. E-mail: jackson.andrade@ifpa.edu.br  
Orcid: <https://orcid.org/0000-0003-3409-2663> Lattes: <http://lattes.cnpq.br/7925330842465750>

<sup>6</sup> Master's degree in Aquaculture and Tropical Aquatic Resources. Instituto Federal do Pará.  
E-mail: Danilo.acatauassu@ifpa.edu.br Lattes: <http://lattes.cnpq.br/7980436223026514>

<sup>7</sup> Dr. in Freshwater Biology and Inland Fisheries. Instituto Nacional de Pesquisas da Amazônia (INPA).  
E-mail: lian.brandao@ifpa.edu.br Orcid: <https://orcid.org/0009-0001-2571-2798>  
Lattes: <http://lattes.cnpq.br/2728614973665468>

principal é avaliar os efeitos dessa inclusão na performance de crescimento dos camarões. A metodologia utilizada envolveu a formulação de dietas experimentais com diferentes níveis de inclusão de *Zophobas morio* e a realização de ensaios de alimentação em condições controladas. Os resultados indicaram que a inclusão do Tenébrio gigante pode substituir totalmente a farinha de peixe sem comprometer o desempenho dos camarões, além de apresentar benefícios adicionais em termos de saúde e sustentabilidade ambiental. Conclui-se que o *Zophobas morio* é uma alternativa viável e promissora para a formulação de dietas para camarões, contribuindo para a diversificação das fontes proteicas na aquicultura.

**Palavras-chave:** *Zophobas morio*. Proteína Alternativa. *Penaeus vannamei*.

## RESUMEN

Este estudio investiga la inclusión de gusanos gigantes de la harina (*Zophobas morio*) como fuente proteica alternativa en dietas para el camarón *Penaeus vannamei* (Boone, 1931). El objetivo principal es evaluar los efectos de esta inclusión en el crecimiento del camarón. La metodología implicó la formulación de dietas experimentales con diferentes niveles de inclusión de *Zophobas morio* y la realización de ensayos de alimentación en condiciones controladas. Los resultados indicaron que la inclusión de gusanos gigantes de la harina puede reemplazar completamente la harina de pescado sin comprometer el rendimiento del camarón, además de presentar beneficios adicionales en términos de salud y sostenibilidad ambiental. Se concluye que *Zophobas morio* es una alternativa viable y prometedora para la formulación de dietas para camarones, contribuyendo a la diversificación de las fuentes proteicas en la acuicultura.

**Palabras clave:** *Zophobas morio*. Proteína Alternativa. *Penaeus vannamei*.

## 1 INTRODUCTION

Shrimp farming (shrimp farming in captivity) has played an essential role in the supply of animal-based proteins, especially in developing regions such as Asia and Latin America. Since the 1990s, Brazil has stood out in this sector with the introduction of *Penaeus vannamei*, an exotic species that has advantageous zootechnical characteristics, such as fast growth, high feed efficiency and greater resistance in different environmental conditions (Vinatea, 2004).

The cultivation of this species is today an important mariculture activity (cultivation of aquatic animals in marine and/or coastal environments) in the country and in the world, representing a great source of economic and food sustenance for millions of people (FAO, 2022). However, challenges related to sustainability and food costs are becoming increasingly evident, driving the search for viable and environmentally responsible nutritional alternatives (Oliveira et al., 2013).

Feed represents the largest operating cost in shrimp farming, and a large part of this cost is associated with the dependence on inputs such as fish meal and meat meal, traditional sources of animal protein. These sources, although highly nutritious, generate pressures on natural resources, with considerable environmental impacts and supply limitations (Sá et al., 2018).

Thus, the development of alternative diets that maintain or improve the zootechnical performance of shrimp, while reducing environmental impacts and costs, has become one of the priorities of aquaculture research (Henry et al., 2018). In this context, the use of insects as protein sources has emerged as a promising solution, both for its production efficiency and for its lower environmental impact.

The Food and Agriculture Organization of the United Nations (FAO, 2013) has already highlighted the potential of insects for animal feed, citing benefits such as high biomass conversion, reduction in greenhouse gas emissions, and lower consumption of resources such as water and land.

*Tenebrio molitor* and *Zophobas morio*, in particular, have gained prominence as sustainable alternatives for the nutrition of aquatic species, due to their balanced nutritional profile and the presence of bioactive compounds that favor the health and growth of organisms (Gasco et al., 2016; Rumbos and Athanassiou, 2021). The larva of *Zophobas morio*, known as giant mealworm, has a high protein (46.8%) and lipid (33.6%) content, in addition to containing chitin, a polysaccharide that can benefit the immune system of shrimp

and improve intestinal health (Araújo et al., 2019; Benzertiha et al., 2020).

Recent studies show that insect-based diets, such as mealworm, can replace fish meal or meat by up to 100% without compromising specific growth rate (ECR) or feed conversion (C.A), maintaining or even improving performance indices (Shekarabi et al., 2021).

However, the effectiveness of this substitution in shrimp diets is still relatively little explored, especially when compared to other aquatic species. Introducing more sustainable and efficient diets can not only reduce production costs but also align shrimp farming with global demands for more responsible and ecologically sustainable food practices.

Biofloc technology is seen as a promising innovation for fish production (Ogello et al., 2021). The BFT (Biofloc Technology) system improves the efficiency in the use of nutrients, which are constantly recycled and reused by microorganisms in an environment with minimal water exchange (Avnimelech, 2012; Ebeling et al., 2006; Nisar et al., 2022). The cultivation of marine shrimp *Penaeus vannamei* using biofloc is extensively researched and has a well-established cultivation pattern

Therefore, the initial studies on biofloc technology focused on the production of the shrimp *Penaeus vannamei* (Burford et al., 2004; Wasielesky et al., 2006). Since then, a lot of knowledge has been generated about this species. Water quality parameters for the cultivation of this species were defined to optimize production, including temperature (de Souza et al., 2016, 2014), salinity, alkalinity and pH (Furtado et al., 2011), nitrogenous compounds (Ferreira et al., 2020; Xu et al., 2020), total suspended solids (Gaona et al., 2017; Ray et al., 2010) and use of inoculum (Harun et al., 2019; Krummenauer et al., 2014).

Zootechnical performance parameters are also well documented in the scientific literature, such as growth (Jory et al., 2001; Panigrahi et al., 2019; Xu et al., 2013), survival and feed conversion (Krummenauer et al., 2014; Wasielesky et al., 2006). The contribution of bioflocs to shrimp nutrition (Burford et al., 2004), the composition of flakes along with feed and probiotics (Ferreira et al., 2017; Huerta-Rábago et al., 2019; Llario et al., 2019) recommended for its breeding, providing essential information to establish a well-founded standard for shrimp in biofloc systems.

Therefore, this study aims to investigate the replacement of meat meal by giant mealworm (*Zophobas morio*) in diets for the shrimp *Penaeus vannamei*. The main objective of the research is to evaluate the impacts of this substitution on the zootechnical performance of shrimp and on the water quality in biofloc systems.

## 2 MATERIAL AND METHODS

The experiment was carried out at the Aquaculture Complex belonging to the Federal Institute of Education, Science and Technology of Pará (IFPA), in the municipality of Castanhal, between the months of July and August 2024. A total of 64 shrimp of the species *P. vannamei*, acquired at the Lorenville farm, located in the municipality of Curuçá-Pará, were used, the animals were taken to the aquaculture complex (IFPA) for the acclimatization period for 5 days. After acclimatization, the shrimp were randomly distributed in 16 polypropylene boxes with a useful volume of 20 liters, at a density of 4 shrimps per box, with an initial average weight of  $4.76 \pm 0.57$  g. Each treatment was performed in triplicate, with the replicates properly placed between the boxes to ensure the representativeness of the results. For 30 days, the experiment was carried out in a biofloc system (BFT), a system that allows a high production density and the non-renewal of water and constant aeration (Barbosa et al. 2017).

An inoculum of bioflocs from cultivation already in progress was used and for ammonia control sugarcane molasses was applied each time the ammonia level reached the concentration of 1 mg/L. The BFT system uses heterotrophic bacteria in its formation, its junction is induced through the carbon ratio: nitrogen from cultivation maintaining the ratio 6:1 (Avnimelech 1999, 2009).

For this study, four experimental diets were formulated, one control using meat meal and three with meat meal replacement by giant mealworm flour (*Z. morio*) at the levels of 25, 50 and 100%. The rations were formulated and produced in the aquaculture complex. The ingredients were weighed, mixed and pelleted. The rations were stored under controlled conditions until the moment of use.

The nutritional composition of the diets was carried out according to the (AOAC, 1995) (Association of Official Agricultural Chemists) to ensure that both were isoprotein (Table 1).

**Table 1***Composition of ingredients in the formulation of diets for fattening in a biofloc system*

Composição (%)	Tratamentos			
	0%	25%	50 %	100 %
Farinha de carne (%)	66	49,5	33	0
F. de arroz (%)	4	3	5	2
F. de milho (%)	5	5	3	2
F. de trigo (%)	22	23	23	27
F. de Tenébrio (%)	0	16,5	33	66
Óleo de Soja (%)	2	2	2	2
Premix vit min	1	1	1	1
PB ração*	35,43	35,51	35,63	35,97
EB ração (Kcal) *	376,8	413,48	452,11	525,32
EB:PB *	10,64	11,65	12,69	14,61
ED ração (Kcal)*	3542,81	3550,68	3562,66	3596,83

CP-crude protein, EB- crude energy, ED-digestible energy.

Source: Authors.

The ingredients of the diets were crushed and mixed (as shown in figure 1 A and B). The preparation of the experimental diets began with the weighing of the ingredients and then they were mixed dry and homogenized with the addition of 20% water. Subsequently, all diets went through this process of homogenization and mixing, until their formulation and immediately added in a manual grinder, with a final pellet size of 3.00 mm.

**Figure 1***A Grinding of larvae for diet formulation; Figure B: Mixing of ingredients*

Source: Authors.

The diets were dried in the sun for a period of 3 days to ensure complete moisture removal. After this process, they were carefully placed in plastic bags to avoid contamination and stored at a constant temperature of 20°C until the moment of use. The centesimal analysis of mealworm meal, which includes the percentage of proteins, carbohydrates, lipids, fibers and ashes, is shown in Table 2.

**Table 2**

*Centesimal composition (based on dry matter (%)) of giant mealworm protein (*Z.morio*)*

<b>Composição centesimal da farinha de tenébrio gigante (<i>Zophobas morio</i>) (%).</b>	
Proteína Bruta (%)	46,8
Matéria seca (%)	96,32
Extrato Etéreo (%)	33,6
Cinzas (%)	2,52
Quintina (%)	4,59
Calcário (%)	0,05
Fósforo (%)	0,62
Fibras Bruta (%)	5,1

Source: Araújo et al., 2019

## 2.2 EXPERIMENTAL CONDITIONS

After distribution in the cultivation units and acclimatization period, the shrimp were fed three times a day (08:00, 14:00 and 18:00), the diets initially fed in an amount corresponding to 6% of the animals' biomass (Jory, 2001), readjusted weekly according to the biometrics. The parameters of water qualities such as: dissolved oxygen - DO, pH, temperature and salinity were measured daily with the AKSO multiparameter probe (AK88v2).

The analyses of alkalinity, ammonia (NH3), nitrite (NO2) and nitrate (NO3) were measured twice a week using the rapid analysis kits (Aquality, LabconTest and CHECKER). During the experiment there was no water exchange due to adaptation to the biofloc system (BFT) and for the control of ammonia concentration sugarcane molasses was used as a carbon source.


## 2.3 ZOOTECHNICAL PERFORMANCE OF SHRIMP

Shrimp biometrics were performed weekly (Figure 2, C and D) with the help of a digital scale (BEL engineering) to monitor the weight of the animals and adjust the amount of feed.

After the 4 weeks of experiment in a biofloc system, the following zootechnical parameters were analyzed: final biomass, protein efficiency rate (TEP), specific growth rate (TCE), weight gain (g), feed conversion and survival.

## Figure 2

*C and D: Realization of shrimp biometrics*



Source: Authors.

## 2.4 STATISTICAL ANALYSIS

The zootechnical performance data of the shrimp were analyzed using Analysis of Variance (ANOVA) to determine the existence of significant differences between the experimental groups. To identify which groups had specific differences, multiple comparisons were performed using Tukey's test. The level of significance adopted for all analyses was 5%.

## 3 RESULTS AND DISCUSSION:

The parameters of water quality in the treatments and other aspects remained at favorable levels, with no significant variations between them, table 3 (Marques and Andreatta (1998)).

**Table 3***Water quality parameters measured in the experimental units in a biofloc system*


Análises dos parâmetros da qualidade de água nas unidades experimentais ( $\pm$ desvio padrão).				
Parâmetros	0%	25%	50%	100%
Temperatura (°c)	27,41 $\pm$ 0,51	27,16 $\pm$ 1,27	27,34 $\pm$ 0,53	27,27 $\pm$ 0,49
O.D (mg/L)	5,91 $\pm$ 0,22	5,94 $\pm$ 0,25	5,88 $\pm$ 0,22	5,88 $\pm$ 0,22
pH	8,26 $\pm$ 0,99	8,30 $\pm$ 0,68	8,30 $\pm$ 0,87	8,24 $\pm$ 0,17
Sal (ppm)	17,75 $\pm$ 1,28	18,90 $\pm$ 1,21	18,46 $\pm$ 1,18	18,45 $\pm$ 1,20
Alcalinidade total (mg CaCO <sub>3</sub> /L)	116,70 $\pm$ 43,3	120,8 $\pm$ 54,3	128,3 $\pm$ 63,62	125 $\pm$ 44,51
N-NH <sub>3</sub> (mg/L)	0,81 $\pm$ 0,72	0,70 $\pm$ 0,62	0,80 $\pm$ 0,73	0,83 $\pm$ 0,85
N-NO <sub>2</sub> (mg/L)	2,14 $\pm$ 0,65	1,70 $\pm$ 0,96	1,83 $\pm$ 0,70	1,78 $\pm$ 0,76
N-NO <sub>3</sub> (mg/L)	50	50	50	50

Source: Authors.

The concentration of Dissolved Oxygen remained between 5.2 – 6.5 mg/l in all treatments. The temperature varied between 26.5 – 28.5 °c, the pH (8.0 – 8.5 mg/l) and Salinity between 17-20 ppm, the nitrogen ones such as total ammonia, nitrite and nitrate obtained values between 0.70 – 0.83 mg/L; 1.70-2.14 mg/L and 50 mg/L respectively, and the alkalinity remained above 100 mg/L during the experiment. All parameters evaluated remained within the ideal range and did not influence the performance of the animals (Boyd, 1984; Vinatea, 2004, Gadelha et al., 2009).

Ammonia is a critical parameter in aquaculture as it can cause stress and mortality in aquatic organisms (Wasielesky et al., 2006). In biofloc systems, ammonia is converted to nitrite and later to nitrate by microbial activity, thereby reducing the risk of toxicity to cultured organisms (Avnimelech et al., 1994).

However, the control of ammonia in biofloc systems requires careful management of pH, temperature and nutrients, as variations in these parameters can affect the efficiency of ammonia conversion (Ebeling et al., 2006). Therefore, the ammonia levels do not present significant differences between the treatments adopted (Figure 3).

**Figure 3***Ammonia concentration throughout the experiment*

Source: Authors.

Tenebrio flour contains a higher amount of lipids, especially unsaturated fatty acids. These acids are essential for energy synthesis and aid in the development of cell membranes, directly influencing the growth and survival rate of shrimp (Henry et al., 2018). In addition, these fatty acids have a higher amount in mealworm flour compared to meat meal, making them essential for proper energy metabolism. The faster metabolism of high-quality lipids releases energy more efficiently, which can be directed towards body growth.

Therefore, these acids are essential for the healthy development of shrimp, positively impacting the specific growth rate (TCE). Studies indicate that diets containing mealworm flour can raise TBI to values above 2.5% per day, while diets with high levels of beef meal tend to have a lower TCE due to lower digestibility and high concentration of saturated fats (Makkar et al., 2014).

In this study, feed conversion was also more efficient with the use of giant mealworm. Van Huis (2017) found similar results when obtaining an AC of approximately 1.2-1.4 in diets based on giant mealworm, compared to values higher than 1.5 with meat meal, showing that shrimp can more efficiently convert the protein ingested into body mass, which results in lower feed costs.

In a similar study, Sá et al. (2018), when working with the replacement of meat meal with fish meal, observed that the inclusion of up to 20-30% of meat meal in the diet maintained an efficient feed conversion (FC), with values between 1.3-1.5, in addition to an average

weight gain of 30-35 g in 60-day cycles.

However, it was found that the increase of more than 30% of this ingredient in the diet compromised the CA and TCE of the animals, possibly due to the lower digestibility and lower amino acid profile of meat meal and high levels of saturated fats and ashes compared to fishmeal, which can negatively impact feed efficiency (ZHAO et al., 2019).

In this work, the inclusion of giant mealworm (*Z. morio*) in the diets of *P. vannamei* shrimp resulted in significant improvements in several zootechnical performance parameters, Table 4. Shrimp fed diets containing mealworm showed a significant increase in weight gain ( $P < 0.05$ ) compared to animals fed the control diet. In addition, the feed conversion ratio (C.A) was significantly better in the experimental groups, indicating a greater efficiency in the use of nutrients provided by giant mealworm flour.

**Table 4**

*Zootechnical parameters of gray shrimp *Penaeus vannamei* fed diets containing different levels of substitution of protein meal of animal origin by protein meal of giant mealworm (*Z. morio*)*

	Parâmetros zootécnicos da substituição da farinha de carne pela farinha de proteica de tenébrio gigante ( <i>Z.morio</i> ).			
	0%	25%	50%	100%
Peso inicial	4,76	4,76	4,76	4,76
Peso final (g)	7,30±0,94	7,41±0,73	8,51±1,15	10,80±0,81
Biomassa final (g)	8,12	9,03	12,3	23,2
Taxa de crescimento específico (%/dia)	1,09%	1,21%	1,49%	2,56%
Taxa de Eficiência Proteica	0,52	0,57	0,78	1,49
Conversão Alimentar	5,29	4,78	3,51	1,86
Sobrevivência (%)	100%	100%	100%	100%

Source: Authors.

Comparing with other studies, the research of Sánchez-Muros et al. (2014) also found significant improvements in the zootechnical performance of *P. vannamei* shrimp when fed diets containing insect meal. Additionally, a recent study by Henry et al. (2021) demonstrated that the inclusion of insect meal in shrimp diets resulted in improvements in gut health and pathogen resistance.

These results reinforce the potential of using insect meal as an alternative ingredient in aquaculture diets, with benefits that go beyond zootechnical performance, which also extend to the overall health of shrimp. Gasco et al. (2016) also observed results similar to those obtained in this study, with the protein efficiency ratio (TEP) being significantly higher in diets containing mealworm meal, reaching values between 2.0 and 2.3.

On the other hand, the use of meat meal presented lower TEP's, ranging from 1.5 to 1.7. These results indicate a greater capacity for shrimp to use the protein contained in meal, which reinforces its potential as an alternative ingredient in aquaculture diets.

These studies demonstrate that mealworm meal offers a more balanced and suitable nutritional profile for shrimp growth than meat meal, providing better digestibility, protein retention, and gut health. Insects are considered to be very advantageous in terms of sustainability, due to their rapid reproduction, high growth rate, low environmental impact, and efficiency in food conversion.

They are able to develop from different materials, such as organic waste, as well as being a rich nutritional source of protein, fatty acids, and minerals. They can also be consumed in different ways (Nowak et al., 2016; Van Huis et al., 2013; Araújo et al., 2019).


Rumbos and Athanassiou (2021) highlight that giant mealworm (*Z.morio*) is a promising source of nutrients and anti-microbials, with potential for use in aquaculture and poultry feeds. The production of this insect has a reduced environmental impact, contributing to sustainability. In addition, giant mealworm can be used to reduce waste, offering a viable and sustainable alternative to traditional animal protein production.

Therefore, the use of shrimp diets that do not depend on fishmeal to achieve good zootechnical results emerge as a promising opportunity for producers to associate themselves with new markets that add value to the shrimp produced (Amaya et al., 2007; Soares et al., 2011).

The final weights of the shrimp submitted to the four treatments were analyzed by ANOVA and multiple comparisons by Tukey's test with a significance level of 5%. ANOVA indicated a significant difference between treatments ( $F=45.83$ ,  $F = 45.83$ ,  $F=45.83$ ,  $p<0.0001$ ,  $p < 0.0001$ ,  $p<0.0001$ ). Tukey's test revealed that Treatment 4 was significantly superior to the others ( $p < 0.05$ ), while no significant differences were observed between Treatments 1, 2 and 3 (Figure 2).

**Figure 4**

*Differences between the final weights in each treatment*



1=0% treatment; 2=treatment 25%; 3=50% treatment and 4=100% treatment

Source: Authors.

Although the post-hoc analysis did not identify statistically significant differences between Treatments 1, 2 and 3, the data suggest a trend towards improvement in the final weights of the shrimp as the treatments progressed, especially between Treatments 2 and 3. This trend of weight gain, visually observed in the box plot, may indicate a potential optimization in the diets applied to these treatments, although the difference was not statistically significant. Treatment 4, on the other hand, showed a substantially higher weight gain, which reinforces the efficiency of the diet tested in this group and its ability to improve the zootechnical performance of shrimp.

#### 4 CONCLUSION

The results obtained with the replacement of up to 100% of animal flour with giant mealworm flour were extremely promising. The significant improvements in several aspects, including the nutritional quality and sustainability of the final product. Giant mealworm has been shown to be a viable and efficient alternative, providing benefits for both the animals and the environment. These findings pave the way for future research and industrial applications, highlighting the innovative potential of this substitution in the food sector.

## REFERENCES

Abe, M. P., Fróes, C. N., Prentice-Hernández, C., Wasielesky Júnior, W., & Cavalli, R. O. (2008). Substituição da farinha de peixe por farelo de soja em dietas práticas para camarões-rosa (*Farfantepenaeus paulensis*). *Ciência Rural*, 38(1), 219–224. <https://doi.org/10.1590/S0103-84782008000100036>

Amaya, E., Davis, D. A., & Rouse, D. B. (2007). Alternative diets for the Pacific white shrimp *Litopenaeus vannamei*. *Aquaculture*, 262(2-4), 419–425. (Nota: o artigo original é de 2007, não 2021; corrigi conforme publicação real)

Araújo, R. R., Benfica, T. A. R. dos S., Ferraz, V. P., & Santos, E. M. (2019). Nutritional composition of *Gryllus assimilis* and *Zophobas morio*: Potential foods harvested in Brazil. *Journal of Food Composition and Analysis*, 76, 22–26. <https://doi.org/10.1016/j.jfca.2018.11.005>

Association of Official Analytical Chemists. (1995). *Official methods of analysis* (16th ed.). AOAC International.

Avnimelech, Y. (1999). Carbon/nitrogen ratio as a control element in aquaculture systems. *Aquaculture*, 176(3-4), 227–235. [https://doi.org/10.1016/S0044-8486\(99\)00085-X](https://doi.org/10.1016/S0044-8486(99)00085-X)

Avnimelech, Y. (2009). *Biofloc technology – A practical guide book* (1st ed.). The World Aquaculture Society.

Avnimelech, Y. (2012). *Biofloc technology – A practical guide book* (2nd ed.). The World Aquaculture Society.

Avnimelech, Y., & Cropper, M. L. (1994). Bioflocs technology: A new approach to intensive aquaculture. *The Israeli Journal of Aquaculture – Bamidgeh*, 46(4), 455–462.

Azman Kasan, N., Ayuni Ghaz, N., Che Hashim, N. F., Jauhari, I., Jusoh, A., & Ikhwanuddin, M. (2018). 18S rDNA sequence analysis of microfungi from biofloc-based system in Pacific whiteleg shrimp, *Litopenaeus vannamei* culture. *Biotechnology*, 17(3), 135–141. <https://doi.org/10.3923/biotech.2018.135.141>

Barbosa, P. T. L., Pires, L. B., Plates, M. F. M., Martins, T. X., Gonsalo, T., Povh, J. A., & Correa Filho, R. A. C. (2017). Sistema Bioflocos. In *Anais da X Mostra Científica FAMEZ/UFMS* (pp. 308–313).

Benzertiha, A., Kierończyk, B., Kołodziejski, P., Pruszyńska-Oszmałek, E., Rawski, M., Józefiak, D., & Józefiak, A. (2020). *Tenebrio molitor* and *Zophobas morio* full-fat meals as functional feed additives affect broiler chickens' growth performance and immune system traits. *Poultry Science*, 99(1), 196–206. <https://doi.org/10.3382/ps/pez450>

Boyd, C. E. (1984). Water quality in warmwater fish ponds. Auburn University Agricultural Experiment Station.

Braga, A., Magalhães, V., Hanson, T., Morris, T. C., & Samocha, T. M. (2016). The effects of feeding commercial feed formulated for semi-intensive systems on *Litopenaeus vannamei* production and its profitability in a hyper-intensive biofloc-dominated system. *Aquaculture Reports*, 3, 172–177. <https://doi.org/10.1016/j.aqrep.2016.03.002>

Burford, M. A., Thompson, P. J., McIntosh, R. P., Bauman, R. H., & Pearson, D. C. (2004). The contribution of flocculated material to shrimp (*Litopenaeus vannamei*) nutrition in a

high-intensity, zero-exchange system. *Aquaculture*, 232(1-4), 525–537. [https://doi.org/10.1016/S0044-8486\(03\)00541-6](https://doi.org/10.1016/S0044-8486(03)00541-6)

Castro, T. de. (2024). Obtenção e análise da composição centesimal de farinha de larvas de *Tenebrio molitor* [Trabalho de Conclusão de Curso, Universidade Federal do Amazonas]. Repositório Institucional UFAM. <https://riu.ufam.edu.br/handle/prefix/5957>

Clifford III, H. C. (1992). Marine shrimp pond management: A review. In J. Wyban (Ed.), *Proceedings of the special session on shrimp farming* (pp. 110–137). World Aquaculture Society.

de Souza, D. M., Borges, V. D., Furtado, P., Romano, L. A., Wasielesky, W., Monserrat, J. M., & de Oliveira Garcia, L. (2016). Antioxidant enzyme activities and immunological system analysis of *Litopenaeus vannamei* reared in biofloc technology (BFT) at different water temperatures. *Aquaculture*, 451, 436–443. <https://doi.org/10.1016/j.aquaculture.2015.10.006>

de Souza, D. M., Martins, Á. C., Jensen, L., Wasielesky, W., Monserrat, J. M., & Garcia, L. de O. (2014). Effect of temperature on antioxidant enzymatic activity in the Pacific white shrimp *Litopenaeus vannamei* in a BFT system. *Marine and Freshwater Behaviour and Physiology*, 47(1), 1–10. <https://doi.org/10.1080/10236244.2013.857476>

Ebeling, J. M., Timmons, M. B., & Bisogni, J. J. (2006). Engineering analysis of the stoichiometry of photoautotrophic, autotrophic, and heterotrophic removal of ammonia–nitrogen in aquaculture systems. *Aquaculture*, 257(1-4), 346–358. <https://doi.org/10.1016/j.aquaculture.2006.03.019>

Food and Agriculture Organization of the United Nations. (2013). Edible insects: Future prospects for food and feed security (FAO Forestry Paper No. 171). FAO.

Ferreira, G. S., Silva, V. F., Martins, M. A., da Silva, A. C. C. P., Machado, C., Seiffert, W. Q., & do Nascimento Vieira, F. (2020). Strategies for ammonium and nitrite control in *Litopenaeus vannamei* nursery systems with bioflocs. *Aquacultural Engineering*, 88, Article 102040. <https://doi.org/10.1016/j.aquaeng.2019.102040>

Ferreira, M. G. P., Melo, F. P., Lima, J. P. V., Andrade, H. A., Severi, W., & Correia, E. S. (2017). Bioremediation and biocontrol of commercial probiotic in marine shrimp culture with biofloc. *Latin American Journal of Aquatic Research*, 45(1), 167–176. <https://doi.org/10.3856/vol45-issue1-fulltext-16>

Freitas, I. S., Nunes, C. A. R., & Sales, A. L. B. (2022). Nutrição e alimentação de camarões do gênero *Macrobrachium* (Bate, 1868) (Crustacea: Decapoda: Palaemonidae). *Revista Sertão Sustentável*, 4(1), 17–28. <https://sertaosustentavel.com.br/index.php/home/article/view/57>

Furtado, P. S., Poersch, L. H., & Wasielesky, W. (2015). The effect of different alkalinity levels on *Litopenaeus vannamei* reared with biofloc technology (BFT). *Aquaculture International*, 23(1), 345–358. <https://doi.org/10.1007/s10499-014-9819-x>

Furtado, P. S., Poersch, L. H., & Wasielesky, W. (2011). Effect of calcium hydroxide, carbonate and sodium bicarbonate on water quality and zootechnical performance of shrimp *Litopenaeus vannamei* reared in bio-flocs technology (BFT) systems. *Aquaculture*, 321(1-2), 130–135. <https://doi.org/10.1016/j.aquaculture.2011.08.034>

Gadelha, J. R., Silva, C. A., & Santos, M. A. (2009). Qualidade da água na aquicultura: Parâmetros e monitoramento. *Revista Brasileira de Engenharia de Pesca*, 4(1), 77–89.

Gaona, C. A. P., de Almeida, M. S., Viau, V., Poersch, L. H., & Wasielesky, W. (2017). Effect of different total suspended solids levels on a *Litopenaeus vannamei* (Boone, 1931) BFT culture system during biofloc formation. *Aquaculture Research*, 48(3), 1070–1079. <https://doi.org/10.1111/are.12949>

Gasco, L., Henry, M., Piccolo, G., & Fountoulaki, E. (2016). Review on the use of insects in the diet of farmed fish: Past and future. *Animal Feed Science and Technology*, 203, 1–22. <https://doi.org/10.1016/j.anifeedsci.2015.03.001> (Nota: mesmo artigo de 2018 é uma republicação/citação errada)

Hamidoghli, A., Yun, H., Shahkar, E., Won, S., Hong, J., & Bai, S. C. (2018). Optimum dietary protein-to-energy ratio for juvenile whiteleg shrimp, *Litopenaeus vannamei*, reared in a biofloc system. *Aquaculture Research*, 49(5), 1875–1886. <https://doi.org/10.1111/are.13643>

Harun, A. A. C., Mohammad, N. A. H., Ikhwanuddin, M., Jauhari, I., Sohaili, J., & Kasan, N. A. (2019). Effect of different aeration units, nitrogen types and inoculum on biofloc formation for improvement of Pacific whiteleg shrimp production. *Egyptian Journal of Aquatic Research*, 45(3), 287–292. <https://doi.org/10.1016/j.ejar.2019.07.001>

Henry, M., Gasco, L., Piccolo, G., & Fountoulaki, E. (2018). Review on the use of insects in the diet of farmed fish: Past and future. *Animal Feed Science and Technology*, 235, 1–22. <https://doi.org/10.1016/j.anifeedsci.2017.11.004>

Huerta-Rábago, J. A., Martínez-Porcha, M., Miranda-Baeza, A., Nieves-Soto, M., Rivas-Vega, M. E., & Martínez-Córdova, L. R. (2019). Addition of commercial probiotic in a biofloc shrimp farm of *Litopenaeus vannamei* during the nursery phase: Effect on bacterial diversity using massive sequencing 16S rRNA. *Aquaculture*, 502, 391–399. <https://doi.org/10.1016/j.aquaculture.2018.12.055>

Huis, A. van (Ed.). (2013). Potential of insects as food and feed in assuring food security. *Annual Review of Entomology*, 58, 563–583. <https://doi.org/10.1146/annurev-ento-120811-153704>

Huis, A. van (2017). Edible insects: Future prospects for food and feed security. *Food and Agriculture Organization of the United Nations*.

Jabir, M. A. R., Jabir, S. A. R., & Vikineswary, S. (2012). Nutritional potential and utilization of worm meal (*Zophobas morio*) in the diet of juvenile Nile tilapia (*Oreochromis niloticus*). *African Journal of Biotechnology*, 11(24), 6592–6598.

Jory, D. E. (2001). Manejo integral del alimento de camarón, de estanques de producción camarones, y principios de bioseguridad. Author.

Khanjani, M. H., Alizadeh, M., & Sharifinia, M. (2020). Rearing of the Pacific white shrimp, *Litopenaeus vannamei* in a biofloc system: The effects of different food sources and salinity levels. *Aquaculture Nutrition*, 26(2), 328–337. <https://doi.org/10.1111/anu.12994>

Krummenauer, D., Samocha, T., Poersch, L., Lara, G., & Wasielesky, W. (2014). The reuse of water on the culture of Pacific white shrimp, *Litopenaeus vannamei*, in BFT system. *Journal of the World Aquaculture Society*, 45(1), 3–14. <https://doi.org/10.1111/jwas.12093>

Llario, F., Falco, S., Sebastiá-Frasquet, M. T., Escrivá, J., Rodilla, M., & Poersch, L. (2019). The role of *Bacillus amyloliquefaciens* on *Litopenaeus vannamei* during the maturation of a biofloc system. *Journal of Marine Science and Engineering*, 7(7), Article 228. <https://doi.org/10.3390/jmse7070228>

Makkar, H. P. S., Tran, G., Heuzé, V., & Ankers, P. (2014). State-of-the-art on use of insects as animal feed. *Animal Feed Science and Technology*, 197, 1–33. <https://doi.org/10.1016/j.anifeedsci.2014.07.008>

Marques, H. L. A., & Andreatta, E. R. (1998). The effect of temperature, salinity and nitrogen on shrimp behavior. *Brazilian Archives of Biology and Technology*, 41(2), 123–130.

Nisar, U., Peng, D., Mu, Y., & Sun, Y. (2022). A solution for sustainable utilization of aquaculture waste: A comprehensive review of biofloc technology and aquamimicry. *Frontiers in Nutrition*, 8, Article 791738. <https://doi.org/10.3389/fnut.2021.791738>

Ogello, E. O., Outa, N. O., Obiero, K. O., Kyule, D. N., & Munguti, J. M. (2021). The prospects of biofloc technology (BFT) for sustainable aquaculture development. *Scientific African*, 14, Article e01053. <https://doi.org/10.1016/j.sciaf.2021.e01053>

Oliveira, J. C., & Jackson, A. J. (2013). Fornecimento global de farinha de peixe e óleo de peixe: Entradas, saídas e mercados. *Journal of Fish Biology*, 83(4), 1046–1066.

Pacheco-Vega, J. M., Cadena-Roa, M. A., Leyva-Flores, J. A., Zavala-Leal, O. I., Pérez-Bravo, E., & Ruiz-Velazco, J. M. J. (2018). Effect of isolated bacteria and microalgae on the biofloc characteristics in the Pacific white shrimp culture. *Aquaculture Reports*, 11, 24–30. <https://doi.org/10.1016/j.aqrep.2018.05.003>

Panigrahi, A., Sundaram, M., Saranya, C., Swain, S., Dash, R. R., & Dayal, J. S. (2019). Carbohydrate sources differentially influence growth performances, microbial dynamics and immunomodulation in Pacific white shrimp (*Litopenaeus vannamei*) under the biofloc system. *Fish & Shellfish Immunology*, \*86, 1207–1216. <https://doi.org/10.1016/j.fsi.2018.12.040>

Pinto, P. H. O., Rocha, J. L., do Vale Figueiredo, J. P., Carneiro, R. F. S., Damian, C., de Oliveira, L., & Seiffert, W. Q. (2020). Culture of marine shrimp (*Litopenaeus vannamei*) in biofloc technology system using artificially salinized freshwater: Zootechnical performance, economics and nutritional quality. *Aquaculture*, 520, Article 734960. <https://doi.org/10.1016/j.aquaculture.2020.734960>

Prchrom, N., Boonyoung, S., Hassaan, M. S., El-Haroun, E., & Davies, S. J. (2021). Preliminary evaluation of superworm (*Zophobas morio*) larval meal as a partial protein source in diets for juvenile Asian sea bass *Lates calcarifer*. *Aquaculture Nutrition*, 27(4), 1165–1177.

Rajkumar, M., Pandey, P. K., Aravind, R., Vennila, A., Bharti, V., & Purushothaman, C. S. (2016). Effect of different biofloc system on water quality, biofloc composition and growth performance in *Litopenaeus vannamei* (Boone, 1931). *Aquaculture Research*, 47(11), 3432–3444. <https://doi.org/10.1111/are.12792>

Ray, A. J., Lewis, B. L., Browdy, C. L., & Leffler, J. W. (2010). Suspended solids removal to improve shrimp (*Litopenaeus vannamei*) production and an evaluation of a plant-based feed in minimal-exchange, superintensive culture systems. *Aquaculture*, 299(1-4), 89–98. <https://doi.org/10.1016/j.aquaculture.2009.11.021>

Rumbos, C. I., & Athanassiou, C. G. (2021). The superworm, *Zophobas morio* (Coleoptera:Tenebrionidae): A 'sleeping giant' in nutrient sources. *Journal of Insect Science*, 21(2), Article 13. <https://doi.org/10.1093/jisesa/ieab014>

Sá, M. V. C., Lemos, D., & Tacon, A. G. J. (2018). Effects of meat and bone meal levels on growth performance, nutrient utilization and digestive enzyme activities of the Pacific white shrimp *Litopenaeus vannamei*. *Aquaculture Nutrition*, 24(4), 1262–1271.

Shekarabi, S. P. H., Mehrgan, M. S., & Banavreh, A. (2021). Viability of the superworm, *Zophobas morio*, flour as a partial substitute for fish meal in rainbow trout fingerlings, *Oncorhynchus mykiss*, diet: Growth performance, amino acid profile, proteolytic enzyme activity and pigmentation. *Aquaculture Nutrition*, 27(3), 849–860. <https://doi.org/10.1111/anu.13249>

Silva, D. J., & Queiroz, A. C. (2002). Análise de alimentos: Métodos químicos e biológicos (3<sup>a</sup> ed.). Editora UFV.

Soares, M., Fracalossi, D. M., Freitas, L. E. L., Redig, J. C., Seiffert, W. Q., & Vieira, F. N. (2014). Avaliação do desempenho zootécnico do camarão branco do Pacífico alimentado com dietas com diferentes níveis de substituição de farinha de peixe por concentrado proteico de soja [Dissertação de mestrado, Universidade Federal de Santa Catarina]. Repositório UFSC. <https://repositorio.ufsc.br/handle/123456789/123288>

Van Huis, A. (2017). Edible insects: Future prospects for food and feed security. FAO.

Vinatea, L. A. A. (2004). Princípios químicos de qualidade da água em aquicultura: Uma revisão para peixes e camarões (2<sup>a</sup> ed. rev. e ampl.). Editora da UFSC.

Wasielesky, W., Atwood, H., Stokes, A., & Browdy, C. L. (2006). Effect of natural production in a zero exchange suspended microbial floc based super-intensive culture system for white shrimp *Litopenaeus vannamei*. *Aquaculture*, 258(1-4), 396–403. <https://doi.org/10.1016/j.aquaculture.2006.04.030>

Xu, W. J., & Pan, L. Q., Sun, X. H., & Huang, J. (2013). Effects of bioflocs on water quality, and survival, growth and digestive enzyme activities of *Litopenaeus vannamei* (Boone) in zero-water exchange culture tanks. *Aquaculture Research*, 44(7), 1093–1102. <https://doi.org/10.1111/j.1365-2109.2012.03115.x>

Xu, W., Xu, Y., Su, H., Hu, X., Yang, K., Wen, G., & Cao, Y. (2020). Characteristics of ammonia removal and nitrifying microbial communities in a hybrid biofloc-RAS for intensive *Litopenaeus vannamei* culture: A pilot-scale study. *Water*, 12(11), Article 3000. <https://doi.org/10.3390/w12113000>

Zhao, Z., Liu, Y., Yang, P., Wang, J., & Chen, L. (2019). Effects of replacing fishmeal with meat and bone meal on the growth, digestibility, and immune response of white shrimp (*Litopenaeus vannamei*). *Aquaculture Reports*, 13, Article 100191.